antibodies -online.com





anti-FRAT2 antibody (AA 131-230) (FITC)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL
Target:	FRAT2
Binding Specificity:	AA 131-230
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FRAT2 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FRAT2	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep	
Purification:	Purified by Protein A.	

Target Details

Target:	FRAT2
Alternative Name:	FRAT2 (FRAT2 Products)

Target Details

Background:	Synonyms: Frequently rearranged in advanced T-cell lymphomas 2 antibodyGSK-3 binding	
	protein FRAT2, FRAT2_HUMAN.	
	Background: FRAT1 and FRAT2 were originally characterized as proteins frequently rearranged	
	in advanced T cell lymphoma, and they have since been identified as proto-oncogenes involved	
	in tumorigenesis. These proteins share significant homology with the Xenopus glycogen	
	synthase kinase-3 (xGSK-3) binding protein, which is designated GBP and is essential for the	
	formation of the dorsal-ventral axis during embryonic development. Establishment of these	
	embryonic axes is mediated by the Wnt intracellular signaling pathway. Wnt signaling is	
	regulated in part by the activity of GSK-3, which phosphorylates and thereby facilitates the	
	degradation of ?catenin. GBP binds to GSK-3 and inhibits this phosphorylation, resulting in the	
	accumulation of ?catenin and the subsequent transcription of Wnt target genes. Like GBP,	
	FRAT2 has been shown to bind xGSK-3, suggesting that FRAT1 and FRAT2 may be GSK-3	
	regulatory proteins.	
Gene ID:	23401	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	