antibodies -online.com





anti-NERP-2 antibody (AA 301-400) (FITC)



()	1/0	r\ /1	014	
()	ve	I V I	-v	V

Quantity:	100 μL	
Target:	NERP-2	
Binding Specificity:	AA 301-400	
Reactivity:	Zebrafish (Danio rerio)	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This NERP-2 antibody is conjugated to FITC	
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NERP2
Isotype:	IgG
Cross-Reactivity:	Zebrafish (Danio rerio)
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NERP-2
Alternative Name:	VGF/NERP2 (NERP-2 Products)

Target Details

Background:	Synonyms: NERP-2, Nerve growth factor inducible, Neuroendocrine regulatory peptide-2, Neurosecretory protein VGF, VGF, VGF_HUMAN. Background: This gene is specifically expressed in a subpopulation of neuroendocrine cells, and is upregulated by nerve growth factor. The structural organization of this gene is similar to that of the rat gene, and both the translated and the untranslated regions show a high degree of sequence similarity to the rat gene. The encoded secretory protein also shares similarities with the secretogranin/chromogranin family, however, its exact function is not known. Nerve growth factor (NGF) is a peptide that plays a key role in the differentiation and survival of neurons in the peripheral nervous system (PNS) and the central nervous system (CNS). VGF is a peptide
	synthesized and secreted by neurons and is upregulated by NGF in the PC12 cell line. VGF is widely expressed in both the PNS and CNS, but is especially abundant in the adult hypothalamus. VGF plays an essential role in how the brain regulates energy expenditure and
	body weight. Its expression is rapidly induced by injury, the circadian clock, and neuronal activity.
Gene ID:	7425
UniProt:	015240
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months