antibodies .- online.com





anti-VNN2 antibody (AA 101-180) (FITC)



| _ | | | | | |
|---|---|----|----|---|---|
| U | V | er | VI | е | W |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | VNN2 |
| Binding Specificity: | AA 101-180 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This VNN2 antibody is conjugated to FITC |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from human VNN2 |
|-----------------------|--|
| Isotype: | IgG |
| Predicted Reactivity: | Human, Mouse, Rat, Dog, Cow, Sheep, Horse, Rabbit |
| Purification: | Purified by Protein A. |

Target Details

| Target: | VNN2 |
|-------------------|--|
| Alternative Name: | VNN2 (VNN2 Products) |
| Background: | Synonyms: FOAP 4, FOAP-4, Glycosylphosphatidyl inositol-anchored protein GPI-80, GPI 80, |

| GPI-80, OTTHUMP00000017228, OTTHUMP00000233789, OTTHUMP00000233791, |
|---|
| pantetheinase, Protein FOAP-4, vanin 2, Vanin-2, Vannin 2, Vascular non-inflammatory molecule |
| 2, VNN2_HUMAN. |

Background: Vanin-2 is a 520 amino acid GPI-anchor protein that belongs to the CN hydrolase family and BTD/VNN subfamily. Widely expressed with high expression in spleen and blood, vanin-2 is suggested to be involved in thymus homing of bone marrow cells, regulation of Integrin 2-mediated cell adhesion, and migration and motility of neutrophils. Vanin-2 exists as five alternatively spliced isoforms and is encoded by a gene mapping to human chromosome 6q23.2.

Gene ID: 8875

UniProt: 095498

Application Details

| Application Notes: | IF(IHC-P) 1:50-200 |
|--------------------|--------------------|
| | IF(IHC-F) 1:50-200 |
| | IF(ICC) 1:50-200 |

Restrictions: For Research Use only

Handling

| Format: | Liquid |
|--------------------|--|
| Concentration: | 1 μg/μL |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol. |
| Preservative: | ProClin |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | -20 °C |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
| Expiry Date: | 12 months |