# antibodies -online.com





## anti-Chromosome 15 Open Reading Frame 58 (C15orf58) (AA 301-385) antibody (FITC)



Go to Product page

( )	1 /	0	rv	/ 1 /	71	Α.
	1//	$\vdash$	1 \/	16		1/1/
$\sim$	v	$\sim$	1 V	١,	_	v v

Quantity:	100 μL	
Target:	Chromosome 15 Open Reading Frame 58 (C15orf58)	
Binding Specificity:	AA 301-385	
Reactivity:	Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	FITC	
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GDPGP1/C15orf58	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Pig,Horse	
Purification:	Purified by Protein A.	

#### **Target Details**

Target:	Chromosome 15 Open Reading Frame 58 (C15orf58)
Alternative Name:	GDPGP1/C15orf58 (C15orf58 Products)

### **Target Details**

Background:	Synonyms: C15orf58, Chromosome 15 open reading frame 58, GDP-D-glucose phosphorylase	
	1, GDP-D-glucose phosphorylase C15orf58, gdpgp1, GDPP1_HUMAN, VTC2.	
	Background: Encoding more than 700 genes, chromosome 15 is made up of approximately 106	
	million base pairs and is about 3 % of the human genome. Angelman and Prader-Willi	
	syndromes are associated with loss of function or deletion of genes in the 15q11-q13 region. In	
	the case of Angelman syndrome, this loss is due to inactivity of the maternal 15q11-q13	
	encoded UBE3A gene in the brain by either chromosomal deletion or mutation. In cases of	
	Prader-Willi syndrome, there is a partial or complete deletion of this region from the paternal	
	copy of chromosome 15. Tay-Sachs disease is a lethal disorder associated with mutations of	
	the HEXA gene, which is encoded by chromosome 15. Marfan syndrome is associated with	
	chromosome 15 through the FBN1 gene. The LOC390637 gene product has been provisionally	
	designated LOC390637 pending further characterization.	
Gene ID:	390637	
Application Details		
Application Notes:	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	