# antibodies .- online.com







# anti-GNPDA1 antibody (AA 201-289) (FITC)



_		
()Ver	view	

Quantity:	100 μL
Target:	GNPDA1
Binding Specificity:	AA 201-289
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GNPDA1 antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GNPDA1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Cow,Sheep,Pig
Purification:	Purified by Protein A.

## **Target Details**

Target:	GNPDA1
Alternative Name:	GNPDA1 (GNPDA1 Products)

## **Target Details**

Background:	Synonyms: EC 3.5.99.6, GlcN6P deaminase 1, Glucosamine 6 phosphate deaminase 1,
	Glucosamine 6 phosphate isomerase 1, Glucosamine-6-phosphate deaminase 1, Glucosamine-
	6-phosphate isomerase 1, GNPDA 1, GNPDA1, GNPI, GNPI1_HUMAN, HLN, KIAA0060, Oscillin,
	Putative glucosamine 6 phosphate isomerase.
	Background: During fertilization in mammals, the sperm activates the egg by causing an
	increase in the level of free cytoplasmic calcium concentration. This increased calcium
	concentration induces a characteristic series of oscillations that trigger egg activation and early
	embryo development. A hamster protein named oscillin is thought to be involved in this
	pathway. The enzyme glucosamine-6-phosphate isomerase (GNPI) or deaminase (GNPDA1)
	and the related protein GNPDA2 are the human homologs of hamster oscillin. GNPDA1 and
	GNPDA2 catalyze the conversion of GNP to fructose-6-phosphate and ammonia. Both proteins
	exist as homohexamers and are ubiquitously expressed with highest expression in testis, ovary
	and heart. Three isoforms of GNPDA2 are expressed due to alternative splicing events.
Gene ID:	10007
Application Details	
Application Notes:	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months