



[Go to Product page](#)

Datasheet for ABIN1710792
anti-C4BP antibody (AA 1-100) (FITC)

Overview

Quantity:	100 µL
Target:	C4BP
Binding Specificity:	AA 1-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C4BP antibody is conjugated to FITC
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human C4 binding protein
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Rat,Dog
Purification:	Purified by Protein A.

Target Details

Target:	C4BP
Alternative Name:	C4 Binding Protein (C4BP Products)

Target Details

Background:	<p>Synonyms: C4b binding protein beta chain, C4b-binding protein beta chain, C4BP, C4BPB, C4BPB_HUMAN, Complement component 4 binding protein beta, Complement component 4 binding protein beta chain, Complement component 4-binding protein beta, OTTHUMP00000034288, OTTHUMP00000034289, OTTHUMP00000034375, OTTHUMP0000003474, Proline rich protein, PRP.</p> <p>Background: C4BPB is a member of a superfamily of proteins composed predominantly of tandemly arrayed short consensus repeats of approximately 60 amino acids. A single, unique beta-chain encoded by this gene assembles with seven identical alpha-chains into the predominant isoform of C4b-binding protein, a multimeric protein that controls activation of the complement cascade through the classical pathway. C4b-binding protein has a regulatory role in the coagulation system also, mediated through the beta-chain binding of protein S, a vitamin K-dependent protein that serves as a cofactor of activated protein C. The genes encoding both alpha and beta chains are located adjacent to each other on human chromosome 1 in the regulator of complement activation gene cluster. Alternative splicing gives rise to multiple transcript variants</p>
-------------	---

Application Details

Application Notes:	<p>IF(IHC-P) 1:50-200</p> <p>IF(IHC-F) 1:50-200</p> <p>IF(ICC) 1:50-200</p>
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Handling

Expiry Date: 12 months