antibodies .- online.com





anti-MS4A2 antibody (AA 51-150) (FITC)



Go to Product page

| () | 1/0 | r\ /1 | 014 | |
|-----|-----|-------|-----|---|
| () | ve | I V I | -v | V |
| | | | | |

| Quantity: | 100 μL |
|----------------------|--|
| Target: | MS4A2 |
| Binding Specificity: | AA 51-150 |
| Reactivity: | Human, Mouse |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This MS4A2 antibody is conjugated to FITC |
| Application: | Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)) |

Product Details

| Immunogen: | KLH conjugated synthetic peptide derived from rat MS4A2 |
|-----------------------|---|
| Isotype: | IgG |
| Cross-Reactivity: | Human, Mouse |
| Predicted Reactivity: | Rat |
| Purification: | Purified by Protein A. |

Target Details

| Target: | MS4A2 |
|-------------------|------------------------|
| Alternative Name: | MS4A2 (MS4A2 Products) |

Target Details

| Background: | Synonyms: FCIGA, Ms4a1, Fcer1b, High affinity immunoglobulin epsilon receptor subunit beta, | |
|---------------------|---|--|
| | FcERI, Fc epsilon receptor I beta-chain, IgE Fc receptor subunit beta, Membrane-spanning 4- | |
| | domains subfamily A member 2, Ms4a2, Fce1b | |
| | Background: High affinity receptor that binds to the Fc region of immunoglobulins epsilon. | |
| | Aggregation of FCER1 by multivalent antigens is required for the full mast cell response, | |
| | including the release of preformed mediators (such as histamine) by degranulation and de novo | |
| | production of lipid mediators and cytokines. Also mediates the secretion of important | |
| | lymphokines. Binding of allergen to receptor-bound IgE leads to cell activation and the release | |
| | of mediators responsible for the manifestations of allergy. | |
| Gene ID: | 25316 | |
| UniProt: | P13386 | |
| Pathways: | Fc-epsilon Receptor Signaling Pathway, Regulation of Leukocyte Mediated Immunity, Positive | |
| | Regulation of Immune Effector Process | |
| Application Details | | |
| Application Notes: | IF(IHC-P) 1:50-200 | |
| | IF(IHC-F) 1:50-200 | |
| | IF(ICC) 1:50-200 | |
| Restrictions: | For Research Use only | |
| Handling | | |
| Format: | Liquid | |
| Concentration: | 1 μg/μL | |
| Buffer: | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and | |
| | 50 % Glycerol. | |
| Preservative: | ProClin | |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be | |
| | handled by trained staff only. | |
| Storage: | -20 °C | |
| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. | |
| Expiry Date: | 12 months | |