.-online.com antibodies

Datasheet for ABIN1711032 anti-GPBAR1 antibody (HRP)



Overview

Uverview		
Quantity:	100 μL	
Target:	GPBAR1	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GPBAR1 antibody is conjugated to HRP	
Application:	Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human GPBAR1	
lsotype:	lgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	
Target Details		
Target:	GPBAR1	
Alternative Name:	GPCR TGR5/GPBAR1 (GPBAR1 Products)	
Background:	Synonyms: BG 37, BG37, G protein coupled bile acid receptor 1, G protein coupled bile acid	
	receptor BG 37, G protein coupled bile acid receptor BG37, G-protein coupled bile acid receptor	
	1, G-protein coupled receptor GPCR19, GPBAR 1, GPBAR_HUMAN, GPBAR1, GPCR 19, GPCR,	
	GPCR19, GPR 131, GPR131, hBG 37, hBG37, hGPCR 19, hGPCR19, M BAR, M-BAR, Membrane	

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1711032 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Precaution of Use:

Handling Advice:

Storage:

5			
	bile acid receptor, Membrane type receptor for bile acids, Membrane-type receptor for bile		
	acids, MGC40597, TGR 5, TGR5.		
	Background: The G protein-coupled receptor TGR5 is a 330-amino acid protein that is almost universally expressed in human tissues including heart, skeletal muscle, spleen, kidney, liver,		
	small intestine, placenta, and leukocytes, but not in brain, colon (without mucosa), thymus, or		
	lung. TGR5 is sensitive to bile acids and responds through a significant mechanism that		
	coordinates energy homeostasis. Bile acids activate mitogen-activated protein (MAP) kinase		
	pathways, specifically induce TGR5 internalization, promote an increase of guanosine 5'-0-3-		
	thio-triphosphate binding in membrane fractions, and cause rapid intracellular cAMP		
	production. Bile acids also provoke TGR5 to suppress macrophage functions. TGR5-controlled		
	signaling pathways may be good candidates for drug targets to treat common metabolic		
	diseases, such as obesity, type II diabetes, hyperlipidemia, and atherosclerosis.		
Gene ID:	151306		
Pathways:	WNT Signaling, Hormone Transport, Sensory Perception of Sound		
Application Details			
Application Notes:	WB 1:300-5000		
	IHC-P 1:200-400		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 ar 50 % Glycerol.		
Preservative:	ProClin		

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN1711032 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

handled by trained staff only.

peroxidase.

-20 °C

This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish

lond	lina
land	
10110	

Storage Comment:

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date:

12 months

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 3/3 | Product datasheet for ABIN1711032 | 03/07/2024 | Copyright antibodies-online. All rights reserved.