

Datasheet for ABIN1711214

anti-Allergin-1 antibody (AA 61-160) (HRP)



Go to Product page

_				
	۱۱ / ۱	rv		۱۸/
	' V '	 ı v	Ι.	v v

Quantity:	100 μL	
Target:	Allergin-1 (MILR1)	
Binding Specificity:	AA 61-160	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This Allergin-1 antibody is conjugated to HRP	
Application:	ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))	
Product Details		
Immunogen:	KLH conjugated synthetic poptide derived from human Allergin-1	

Immunogen:	KLH conjugated synthetic peptide derived from human Allergin-1
Isotype:	IgG
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	Allergin-1 (MILR1)	
Alternative Name:	Allergin-1 (MILR1 Products)	
Background: Synonyms: C17orf60,Chromosome 17 open reading frame 60.		

Background: As a major event in type allergic reaction, degranulation of mast cells is triggered by the interaction between specific IgEs and high affinity IgE receptors (FcRIs) on mast cell membrane, and is followed by the release of a battery of inflammatory mediators, such as histamine, leukotrienes and arachidonic acid, and these inflammatory mediators in turn induce a series of allergic symptoms. Recently, Allergin-1 has been found to be an immunoglobulin-like receptor preferentially expressed on mast cells. The crosslinking of Allergin-1 to FcRIs on mast cells may inhibit the signal transduction from FcRIs to intracellular immunoreceptor tyrosine-based inhibitory motif (ITIM), generate ITIM inhibitory signals, which then participate in the regulation of mast cell degranulation, and finally block the initiation of allergic inflammation.

Application Details

Restrictions:	For Research Use only
	IHC-F 1:100-500
Application Notes:	IHC-P 1:200-400

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months