



[Go to Product page](#)

Datasheet for ABIN1711216  
**anti-APLF antibody (AA 421-511) (HRP)**

### Overview

Quantity:	100 µL
Target:	APLF
Binding Specificity:	AA 421-511
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This APLF antibody is conjugated to HRP
Application:	ELISA, Western Blotting (WB), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human APLF
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat
Purification:	Purified by Protein A.

### Target Details

Target:	APLF
Alternative Name:	APLF ( <a href="#">APLF Products</a> )
Background:	Synonyms: 2010301N04Rik, AI452191, Aplf, APLF_HUMAN, Aprataxin and pnk-like factor,

## Target Details

---

Apurinic-apyrimidinic endonuclease APLF, C2orf13, PNK and APTX like FHA protein, PNK and APTX-like FHA domain-containing protein, RGD1565557, XIP1, XRCC1 interacting protein 1, XRCC1-interacting protein 1.

Background: APLF is a 511 amino acid protein that contains one FHA domain and two C2H2-type zinc fingers. Localized to both the nucleus and the cytoplasm, APLF interacts with XRCC1, XRCC4 and Ku-86 and, via these interactions, is involved in single-strand and double-strand DNA break repair. APLF is subject to post-translational phosphorylation in response to DNA breaks. The gene encoding APLF maps to human chromosome 2, which houses over 1,400 genes and comprises nearly 8 % of the human genome. Harlequin ichthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene, while the lipid metabolic disorder sitosterolemia is associated with defects in the ABCG5 and ABCG8 genes. Additionally, an extremely rare recessive genetic disorder, is caused by mutations in the ALMS1 gene, which maps to chromosome 2.

---

Gene ID: 200558

---

Pathways: [DNA Damage Repair](#)

## Application Details

---

Application Notes: WB 1:300-5000  
IHC-P 1:200-400  
IHC-F 1:100-500

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Concentration: 1 µg/µL

---

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

---

Preservative: ProClin

---

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

---

Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

---

## Handling

---

Storage: -20 °C

---

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

Expiry Date: 12 months