# antibodies -online.com





# anti-CYP11B2 antibody (AA 251-350) (HRP)



Go to Product page

$\sim$			
	N/P	r\/I	i⊢₩

Quantity:	100 μL	
Target:	CYP11B2	
Binding Specificity:	AA 251-350	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CYP11B2 antibody is conjugated to HRP	
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human CYP11B2	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Rabbit	
Purification:	Purified by Protein A.	

## Target Details

Target:	CYP11B2
Alternative Name:	CYP11B2 (CYP11B2 Products)

## **Target Details**

Background:	Synonyms: CYP 11B2, CYP11B2, CYPXI11B2, Cytochrome P450 1111B2, Cytochrome P450	
Ç	1111B2 mitochondrial, Cytochrome P450 family 11 subfamily B polypeptide 2, Cytochrome	
	P450 subfamily XIB cholesterol side chain cleavage, Cytochrome P450 subfamily XI11B2,	
	Cytochrome P450C1111B2, C11B2_HUMAN.	
	Background: This gene encodes a member of the cytochrome P450 superfamily of enzymes.	
	The cytochrome P450 proteins are monooxygenases which catalyze many reactions involved	
	in drug metabolism and synthesis of cholesterol, steroids and other lipids. This protein localizes	
	to the mitochondrial inner membrane. The enzyme has steroid 18-hydroxylase activity to	
	synthesize aldosterone and 18-oxocortisol as well as steroid 11 beta-hydroxylase activity.	
	Mutations in this gene cause corticosterone methyl oxidase deficiency. [provided by RefSeq, Jul	
	2008].	
Pathways:	ACE Inhibitor Pathway, Metabolism of Steroid Hormones and Vitamin D, Steroid Hormone	
	Biosynthesis, Regulation of Systemic Arterial Blood Pressure by Hormones, C21-Steroid	
	Hormone Metabolic Process, Feeding Behaviour	
Application Details		
Application Notes:	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

$\vdash$	land	lına
	iaria	шц

Expiry Date:

12 months