antibodies - online.com







anti-DYNC1I2 antibody (AA 61-160) (HRP)



\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	DYNC1I2	
Binding Specificity:	AA 61-160	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This DYNC1I2 antibody is conjugated to HRP	
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DYNC112	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Sheep,Pig,Horse,Rabbit	
Purification:	Purified by Protein A.	

Target Details

Target:	DYNC1I2
Alternative Name:	DYNC1I2 (DYNC1I2 Products)

Target Details

rarget Details	
Background:	Synonyms: Cytoplasmic dynein 1 intermediate chain 2, Cytoplasmic dynein intermediate chain
	2, Dynein intermediate chain 2, cytosolic, DH IC-2, DC1I2_HUMAN, DNCI2, DNCIC2, Dynein
	cytoplasmic intermediate polypeptide 2.
	Background: The inner- and outer-arm dyneins, which bridge between the doublet microtubules
	in axonemes, are the force-generating proteins responsible for the sliding movement in
	axonemes. The intermediate and light chains, thought to form the base of the dynein arm, help
	mediate attachment and may also participate in regulating dynein activity. This gene encodes
	an intermediate chain dynein, belonging to the large family of motor proteins. Mutations in this
	gene result in abnormal ciliary ultrastructure and function associated with primary ciliary
	dyskinesia (PCD) and Kartagener syndrome. [provided by RefSeq, Jul 2008].
Pathways:	M Phase
Application Details	
Application Notes:	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish
	peroxidase.

Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

-20 °C

12 months

Storage:

Expiry Date:

Storage Comment: