



[Go to Product page](#)

Datasheet for ABIN1711310  
**anti-GPR17 antibody (AA 2-100) (HRP)**

### Overview

Quantity:	100 µL
Target:	GPR17
Binding Specificity:	AA 2-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GPR17 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human G protein coupled receptor 17
Isotype:	IgG
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

### Target Details

Target:	GPR17
Alternative Name:	GPR17 ( <a href="#">GPR17 Products</a> )
Background:	Synonyms: DKFZp686M18273, G protein coupled receptor 17, G-protein coupled receptor 17,

## Target Details

---

GPR 17, Gpr17, GPR17\_HUMAN, P2Y like receptor, P2Y-like receptor, Probable P2Y purinoceptor GPR 17, Probable P2Y purinoceptor GPR17, R12, UDP/CysLT receptor, Uracil nucleotide/cysteinyl leukotriene receptor, GPCR GPR17.

Background: G protein-coupled receptor 17, GPR17, also known as uracil nucleotide/cysteinyl leukotriene receptor or P2Y-like receptor (P2YL), is a 367 amino acid member of the G-protein coupled receptor 1 family of proteins. While GPR17 is expressed in kidney, heart and umbilical vein endothelial cells, it is expressed in the highest levels in the brain. Upon brain injury, the extracellular concentrations of nucleotides and cysteinyl leukotrienes (CysLTs), two families of endogenous signaling molecules, increase significantly at the site of damage. In some neurons, GPR17, a membrane receptor for uracil nucleotide and CysLTs, is upregulated as well, infiltrating the lesioned area. GPR17 is thought to play a role in mediating neuronal death, remodeling brain circuitries by microglia and initiating remyelination in damaged neurons. Two named isoforms of GPR17 exist as a result of alternative splicing events.

---

Gene ID: 2840

---

UniProt: [Q13304](#)

## Application Details

---

Application Notes: WB 1:300-5000  
IHC-P 1:200-400  
IHC-F 1:100-500

---

Restrictions: For Research Use only

## Handling

---

Format: Liquid

---

Concentration: 1 µg/µL

---

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

---

Preservative: ProClin

---

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

---

Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

---

## Handling

---

Storage: -20 °C

---

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

---

Expiry Date: 12 months