# antibodies -online.com





# anti-GPR20 antibody (AA 51-150) (HRP)



Go to Product page

( )	11/	IN	/ie	A .
	/ // <del> </del>	۱ ات	/   (−	' \/\/

Quantity:	100 μL	
Target:	GPR20	
Binding Specificity:	AA 51-150	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GPR20 antibody is conjugated to HRP	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

## **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human G protein coupled receptor 20	
Isotype:	IgG	
Cross-Reactivity:	Human	
Purification:	Purified by Protein A.	

# **Target Details**

Target:	GPR20	
Alternative Name:	GPR20 (GPR20 Products)	
Background:	Synonyms: G protein coupled receptor 20, G-protein coupled receptor 20, Gpr20,	

GPR20\_HUMAN, GPCR GPR20.

Background: G protein-coupled receptors (GPRs), also known as seven transmembrane receptors, heptahelical receptors or 7TM receptors, comprise a superfamily of proteins that play a role in many different stimulus-response pathways. GPR signaling is an evolutionarily ancient mechanism used by all eukaryotes to sense environmental stimuli and mediate cell-cell communication. G protein-coupled receptors translate extracellular signals into intracellular signals (G protein activation) and they respond to a variety of signaling molecules, such as hormones and neurotransmitters. GPR20 is a 358 amino acid membrane protein that constitutively activates G(i) proteins without ligand stimulation. Also, GPR20 may be involved in the control of intracellular cAMP levels and mitogenic signaling. Interestingly, GPR20 is expressed in liver and certain regions of the brain, including putamen, caudate and thalamus, but is not expressed in hypothalamus, pons and frontal cortex.

Gene ID:

2843

### **Application Details**

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

	lI	1:
-	เวทก	111111
	land	11110

Expiry Date:

12 months