antibodies -online.com





anti-IQSEC2 antibody (AA 851-950) (HRP)



Go to Product page

\sim					
()	VE	۲۱	/1	\triangle	Λ

Quantity:	100 μL	
Target:	IQSEC2	
Binding Specificity:	AA 851-950	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This IQSEC2 antibody is conjugated to HRP	
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human IQSEC2	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig	
Purification:	Purified by Protein A.	

Target Details

Target:	IQSEC2	
Alternative Name:	IQSEC2 (IQSEC2 Products)	

Target Details

Background:

Synonyms: BRAG1, IQ mot and SEC7 domain-containing protein 2, IQEC2_HUMAN, Iqsec2, KIAA0522, mKIAA0522, MRX1, RP11-258C19.1.

Background: IQSEC2 is a 1,478 amino acid protein that belongs to the BRAG family and contains one IQ domain, one PH domain and a SEC7 domain. Localizing to the cytoplasm, IQSEC2 is expressed in brain, kidney and small intestine, with weaker levels of expression in placenta, pancreas, ovary, prostate and liver. IQSEC2 is a component of the postsynaptic density at excitatory synapses, and interacts with ARF family members as a guanine nucleotide exchange factor. Through the activation of ARF substrates, IQSEC2 may play a crucial role in cytoskeletal and synaptic organization. The gene encoding IQSEC2 maps to the human X chromosome. Defects to the IQSEC2 gene have been linked to mental retardation X-linked type 1 (MRX1), a condition characterized by decreased intellectual function. IQSEC2 exists as three isoforms due to alternative splicing events.

Application Details

Restrictions:	For Research Lise only
	IHC-F 1:100-500
Application Notes:	IHC-P 1:200-400

Handling

- I arialing		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	