

# Datasheet for ABIN1711376

# anti-MAP4K5 antibody (AA 751-846) (HRP)



#### Overview

Overview	
Quantity:	100 μL
Target:	MAP4K5
Binding Specificity:	AA 751-846
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MAP4K5 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)),
	Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human MAP4K5
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Cow,Horse,Chicken
Purification:	Purified by Protein A.
Target Details	
Target Details  Target:	MAP4K5

#### Target Details

Background:
-------------

Synonyms: EC 2.7.11.1, KHS, Kinase homologous to SPS1/STE20, MAPK/ERK kinase kinase kinase 5, MEK kinase kinase 5, MEKKK 5, M4K5\_HUMAN.

Background: Several mammalian kinases have been identified with sequence similarity to the Saccharomyces cerevisiae serine/threonine kinase STE20. STE20 is involved in relaying signals from G-protein coupled receptors to cytosolic MAP kinase cascades, and it lies upstream of a MAP kinase kinase kinase. Mammalian STE20-like kinases include KHS, GLK, NIK, YSK1, HPK1, Krs-1, Krs-2 and GC kinase. KHS (for kinase homologous to SPS1/STE20) is a protein that is

most closely related to GC kinase. The KHS kinase has been shown to activate a variety of

substrates, including JNK, suggesting a role in stress response.

### **Application Details**

Application Notes:

WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months