# antibodies -online.com





### Datasheet for ABIN1711381

## anti-MTERF antibody (AA 101-200) (HRP)



#### Go to Product page

( )	ve	K\ /		A .
	$\cup$	1 V/	Щ.	V۷

Quantity:	100 μL
Target:	MTERF
Binding Specificity:	AA 101-200
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MTERF antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human MTERF1	
Isotype:	IgG	
Predicted Reactivity:	Human,Mouse,Rat	
Purification:	Purified by Protein A.	

#### **Target Details**

Target:	MTERF
Alternative Name:	MTERF1 (MTERF Products)
Background:	Synonyms: mitochondrial, Mitochondrial transcription termination factor 1, MTERF1, mTERF,

MTERF HUMAN.	Transcription	termination factor.

Background: Members of the mTERF family, including MTERF, MTERFD1, MTERFD2 and MTERFD3, are mitochondrial proteins that are believed to be transcription termination factors. MTERF (mitochondrial transcription termination factor 1) is composed of 399 amino acids and contains three leucine zippers that form a three-stranded coiled-coil that binds to DNA. It has been suggested that only the phosphorylated form of MTERF has transcription termination activity. MTERFD1 is also thought to act as a mitochondrial transcription regulator and is expressed as two isoforms produced by alternative splicing. MTERFD3 is believed to be involved in cell cycle regulation and cell growth by modulating mitochondrial transcription. MTERFD3 is expressed in heart, skeletal muscle, pancreas and liver.

Gene ID:

7978

#### **Application Details**

Application Notes:	WB 1:300-5000

IHC-P 1:200-400 IHC-F 1:100-500

Restrictions: For Research Use only

#### Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.
Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months