antibodies -online.com





anti-MYCBPAP antibody (AA 401-500) (HRP)



Go to Product page

()	ve	K\ /		A .
	\cup	1 V/	Щ.	V۷

Quantity:	100 μL
Target:	MYCBPAP
Binding Specificity:	AA 401-500
Reactivity:	Rat, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MYCBPAP antibody is conjugated to HRP
Application:	ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human MYCBPAP
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human
Purification:	Purified by Protein A.

Target Details

Target:	MYCBPAP
Alternative Name:	MYCBPAP (MYCBPAP Products)

Target Details

Bacl	//	ro	ıın	٨.
Daci	ĸα	ΙU	uu	u.

Synonyms: AMAM-1, AMAP-1, AMAP1, AMY-1-binding protein 1, DKFZp434N1415,

MYBPP_HUMAN, MYCBP-associated protein, Mycbpap.

Background: AMAP-1 (AMY-1-binding protein 1), also known as AMAM-1 or MYCBPAP (MYCBP associated protein), is a 947 amino acid protein that is expressed specifically in testis and is involved in spermatogenesis and synaptic processes. AMAP-1 colocalizes with MYCBP (AMY-1) in cytoplasm and also localizes to membrane. The gene encoding AMAP-1 maps to human chromosome 17, which comprises over 2.5 % of the human genome and encodes over 1,200 genes. Two key tumor suppressor genes are associated with chromosome 17, namely, p53 and BRCA1. Tumor suppressor p53 is necessary for maintenance of cellular genetic integrity by moderating cell fate through DNA repair versus cell death. Malfunction or loss of p53 expression is associated with malignant cell growth and Li-Fraumeni syndrome. Like p53, BRCA1 is directly involved in DNA repair, though specifically it is recognized as a genetic determinant of early onset breast cancer and predisposition to cancers of the ovary, colon, prostate gland and fallopian tubes.

Pathways:

Sensory Perception of Sound

Application Details

Application Notes:

IHC-P 1:200-400

IHC-F 1:100-500

-20 °C

Restrictions:

For Research Use only

Handling

Storage:

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months