antibodies - online.com







anti-TUBGCP3 antibody (AA 401-500) (HRP)



\sim			
()	ver	VIA	۱۸/
\sim	v Cı	v 1 C	v v

Quantity:	100 μL	
Target:	TUBGCP3	
Binding Specificity:	AA 401-500	
Reactivity:	Xenopus laevis	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This TUBGCP3 antibody is conjugated to HRP	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human GCP3/Gamma-tubulin complex component 3	
Isotype:	IgG	
Cross-Reactivity:	Xenopus laevis	
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Chicken	
Purification:	Purified by Protein A.	
Target Details		
Target:	TUBGCP3	

Target Details

Alternative Name:	GCP3/Gamma-tubulin complex component 3 (TUBGCP3 Products)	
Background:	Synonyms: Gamma-ring complex protein 104 kDa, Gamma-tubulin complex component 3, GCP	
	3, GCP3, GCP3_HUMAN, h104p, hGCP3, hGrip104, hSpc98, SPBC98, Spc98p, Spindle pole body	
	protein Spc98 homolog, TUBGCP3, Tubulin gamma complex associated protein 3.	
	Background: The Gamma-Tubulin complex is composed of Gamma Tubulin and the Gamma-	
	Tubulin complex-associated proteins GCP2, GCP3, GCP4, GCP5 and GCP6, all of which are	
	essential components of microtubule organizing centers. Gamma-Tubulin complex	
	components are localized to both the centrosome, where they are involved in microtubule	
	nucleation, and to the cytoplasm, where they exist as soluble complexes that can be recruited	
	to the centrosome as needed. Although the GCP proteins are related, they have distinct roles	
	which contribute to the proper function of the Gamma-Tubulin complex. GCP3 (Gamma-	
	Tubulin complex component 3), also known as TUBGCP3 or SPBC98, localizes to the	
	centrosome and is a ubiquitously expressed 907 amino acid member of the Gamma-Tubulin	
	complex. Like GCP2 and Gamma Tubulin, GCP3 is conserved in all eukaryotes, suggesting that	
	it is part of a core unit involved in eukaryotic microtubule nucleation. Three isoforms of GCP3	
	exist due to alternative splicing events.	
Gene ID:	10426	
Pathways:	M Phase, SARS-CoV-2 Protein Interactome	
Application Details		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	

Handling

	handled by trained staff only.	
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	