Datasheet for ABIN1711586 anti-TCTN3 antibody (AA 201-300) (HRP)

-online.com antibodies



$\sim$		
Οv	Pr	۱۸/
$\bigcirc$ v		, v v

Quantity:	100 μL
Target:	TCTN3
Binding Specificity:	AA 201-300
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TCTN3 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

## Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TCTN3/TECT3
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Chicken,Rabbit
Purification:	Purified by Protein A.

## Target Details

Target:	TCTN3
Alternative Name:	TCTN3/TECT3 (TCTN3 Products)
Background:	Synonyms: C10orf61, Chromosome 10 open reading frame 61, DKFZP564D116, TCTN3,

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1711586 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

<ul> <li>Background: Tect3 is a 607 amino acid single-pass type I membrane protein that belongs to the</li> <li>tectonic family and exists as four alternatively spliced isoforms. Tect3 interacts with MKS1 and</li> <li>may be involved in apoptosis regulation. The gene that encodes Tect3 contains approximately</li> <li>31,560 bases and maps to human chromosome 10q24.1. Spanning nearly 135 million base</li> <li>pairs and encoding nearly 1,200 genes, chromosome 10 makes up approximately 4.5 % of the</li> <li>human genome. Several protein-coding genes, including those that encode chemokines,</li> <li>cadherins, excision repair proteins, early growth response factors (Egrs) and fibroblast growth</li> <li>receptors (FGFRs), are located on chromosome 10. Defects in some of the genes that map to</li> <li>chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome,</li> <li>Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrome,</li> <li>Cockayne syndrome, multiple endocrine neoplasia type 2 and porphyria. Tetrahydrobiopterin</li> <li>deficiency and a number of syndromes involving defective skull and facial bone fusion are also</li> <li>linked to chromosome 10.</li> </ul>	TECT3_HUMAN, Tectonic 3, Tectonic 3 precursor, Tectonic family member 3, Tectonic-3.
may be involved in apoptosis regulation. The gene that encodes Tect3 contains approximately 31,560 bases and maps to human chromosome 10q24.1. Spanning nearly 135 million base pairs and encoding nearly 1,200 genes, chromosome 10 makes up approximately 4.5 % of the human genome. Several protein-coding genes, including those that encode chemokines, cadherins, excision repair proteins, early growth response factors (Egrs) and fibroblast growth receptors (FGFRs), are located on chromosome 10. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrome, Cockayne syndrome, multiple endocrine neoplasia type 2 and porphyria. Tetrahydrobiopterin deficiency and a number of syndromes involving defective skull and facial bone fusion are also	Background: Tect3 is a 607 amino acid single-pass type I membrane protein that belongs to the
31,560 bases and maps to human chromosome 10q24.1. Spanning nearly 135 million base pairs and encoding nearly 1,200 genes, chromosome 10 makes up approximately 4.5 % of the human genome. Several protein-coding genes, including those that encode chemokines, cadherins, excision repair proteins, early growth response factors (Egrs) and fibroblast growth receptors (FGFRs), are located on chromosome 10. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrome, Cockayne syndrome, multiple endocrine neoplasia type 2 and porphyria. Tetrahydrobiopterin deficiency and a number of syndromes involving defective skull and facial bone fusion are also	tectonic family and exists as four alternatively spliced isoforms. Tect3 interacts with MKS1 and
pairs and encoding nearly 1,200 genes, chromosome 10 makes up approximately 4.5 % of the human genome. Several protein-coding genes, including those that encode chemokines, cadherins, excision repair proteins, early growth response factors (Egrs) and fibroblast growth receptors (FGFRs), are located on chromosome 10. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrome, Cockayne syndrome, multiple endocrine neoplasia type 2 and porphyria. Tetrahydrobiopterin deficiency and a number of syndromes involving defective skull and facial bone fusion are also	may be involved in apoptosis regulation. The gene that encodes Tect3 contains approximately
human genome. Several protein-coding genes, including those that encode chemokines, cadherins, excision repair proteins, early growth response factors (Egrs) and fibroblast growth receptors (FGFRs), are located on chromosome 10. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrome, Cockayne syndrome, multiple endocrine neoplasia type 2 and porphyria. Tetrahydrobiopterin deficiency and a number of syndromes involving defective skull and facial bone fusion are also	31,560 bases and maps to human chromosome 10q24.1. Spanning nearly 135 million base
cadherins, excision repair proteins, early growth response factors (Egrs) and fibroblast growth receptors (FGFRs), are located on chromosome 10. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrome, Cockayne syndrome, multiple endocrine neoplasia type 2 and porphyria. Tetrahydrobiopterin deficiency and a number of syndromes involving defective skull and facial bone fusion are also	pairs and encoding nearly 1,200 genes, chromosome 10 makes up approximately 4.5 % of the
receptors (FGFRs), are located on chromosome 10. Defects in some of the genes that map to chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrome, Cockayne syndrome, multiple endocrine neoplasia type 2 and porphyria. Tetrahydrobiopterin deficiency and a number of syndromes involving defective skull and facial bone fusion are also	human genome. Several protein-coding genes, including those that encode chemokines,
chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome, Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrome, Cockayne syndrome, multiple endocrine neoplasia type 2 and porphyria. Tetrahydrobiopterin deficiency and a number of syndromes involving defective skull and facial bone fusion are also	cadherins, excision repair proteins, early growth response factors (Egrs) and fibroblast growth
Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrome, Cockayne syndrome, multiple endocrine neoplasia type 2 and porphyria. Tetrahydrobiopterin deficiency and a number of syndromes involving defective skull and facial bone fusion are also	receptors (FGFRs), are located on chromosome 10. Defects in some of the genes that map to
Cockayne syndrome, multiple endocrine neoplasia type 2 and porphyria. Tetrahydrobiopterin deficiency and a number of syndromes involving defective skull and facial bone fusion are also	chromosome 10 are associated with Charcot-Marie Tooth disease, Jackson-Weiss syndrome,
deficiency and a number of syndromes involving defective skull and facial bone fusion are also	Usher syndrome, nonsyndromatic deafness, Wolman?s syndrome, Cowden syndrome,
	Cockayne syndrome, multiple endocrine neoplasia type 2 and porphyria. Tetrahydrobiopterin
linked to chromosome 10.	deficiency and a number of syndromes involving defective skull and facial bone fusion are also
	linked to chromosome 10.

Gene ID:

26123

## Application Details

Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only

## Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN1711586 | 03/07/2024 | Copyright antibodies-online. All rights reserved.

Handling	ndling	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	