

Datasheet for ABIN1711600 anti-CRTAP antibody (AA 201-300) (HRP)



Go to Product page

Overview		
Quantity:	100 μL	
Target:	CRTAP	
Binding Specificity:	AA 201-300	
Reactivity:	Human	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This CRTAP antibody is conjugated to HRP	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human CRTAP	
Isotype:	IgG	
Cross-Reactivity:	Human	
Predicted Reactivity:	Mouse,Rat,Dog,Pig,Horse,Chicken,Rabbit	
Purification:	Purified by Protein A.	
Target Details		
Target:	CRTAP	
Alternative Name:	CRTAP (CRTAP Products)	

Target Details

Background:	Synonyms: Cartilage associated protein, LEPREL3, leprecan-like 3, CASP, CRTAP_HUMAN. Background: CRTAP is a secreted protein localizing to the extracellular space that plays a role in collagen post-translational modifications, extracellular fibril assembly and intracellular trafficking. CRTAP is widely expressed with predominant expression in articular chondrocytes. It contains a signal peptide and a tetratricopeptide-like helical domain and is essential for normal bone formation. In the endoplasmic reticulum (ER), CRTAP forms a complex with Gros1 and CyPB (cyclophilin B) and is required for the efficient 3-hydroxylation of target prolyl residues	
	in Collagen Type I molecules, the major structural proteins of skin and bone. Mutations in the gene encoding CRTAP can lead to autosomal recessive osteogenesis imperfecta (OI) type 7 and type 2B. OI, also known as brittle bone disease, is characterized by bone fragility and	
	susceptibility to fractures. OI type 7 is a mild form of this disorder, while OI type 2B is a neonatal lethal condition.	
Gene ID:	10491	
Application Details		
Application Notes:	WB 1:300-5000 IHC-P 1:200-400 IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	

\vdash	land	lına
	iaria	шц

Expiry Date:

12 months