

Datasheet for ABIN1711669 anti-EXOSC7 antibody (AA 21-120) (HRP)



()	ve	rvi	6	W
\sim	v C	1 V I	\sim	v v

Overview	
Quantity:	100 μL
Target:	EXOSC7
Binding Specificity:	AA 21-120
Reactivity:	Human, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This EXOSC7 antibody is conjugated to HRP
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	

Immunogen:	KLH conjugated synthetic peptide derived from human EXOSC7
Isotype:	IgG
Cross-Reactivity:	Human, Rat
Predicted Reactivity:	Mouse,Dog,Cow,Sheep,Pig,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	EXOSC7
Alternative Name:	EXOSC7 (EXOSC7 Products)

Target Details

Background:	Synonyms: EAP1, EXOS7_HUMAN, Exosc7, Exosome complex component RRP42, Exosome complex exonuclease RRP42, Exosome component 7, FLJ26543, hRrp42p, KIAA0116, p8, Ribosomal RNA processing protein 42, Ribosomal RNA-processing protein 42, RRP42, Rrp42p. Background: The exosome is a multisubunit complex of 3? to 5? exoribonucleases. It is involved in a variety of cellular processes and is responsible for degrading unstable mRNAs that contain AU-rich elements in their untranslated 3? region. EXOSC7 (exosome component 7), also known as p8, EAP1, RRP42 (Ribosomal RNA-processing protein 42), Rrp42p or hRrp42p, is a component of the exosome multienzyme ribonuclease complex. It belongs to the RNase PH family and localizes to the nucleolus. EXOSC7 is one of the six RNase-PH domain subunits of the exosome. Together, these six subunits form a PNPase-like ring. EXOSC7 is required for the processing of the 7S pre-RNA.	
Gene ID:	23016	
Application Details		
Application Notes:	IHC-P 1:200-400 IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.	
Storage:	-20 °C	
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.	
Expiry Date:	12 months	