# antibodies -online.com







## anti-MEGF10 antibody (AA 55-150) (HRP)



Go to Product page

$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL	
Target:	MEGF10	
Binding Specificity:	AA 55-150	
Reactivity:	Mouse	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This MEGF10 antibody is conjugated to HRP	
Application:	ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffinembedded Sections) (IHC (p))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human MEGF10	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken	
Purification:	Purified by Protein A.	

#### Target Details

Target:	MEGF10
Alternative Name:	MEGF10 (MEGF10 Products)

#### **Target Details**

Background:	Synonyms: MEG10_HUMAN, MEGF protein, megf10, Multiple EGF-like domains protein 10,
	multiple EGF-like-domains 10, Multiple epidermal growth factor-like domains 10, Multiple
	epidermal growth factor-like domains protein 10.
	Background: MEGF10 is a 1,140 amino acid protein encoded by the human gene MEGF10.
	MEGF10 belongs to the MEGF family and contains fifteen EGF-like domains and one EMI
	domain. The engulfment of dying cells is a specialized form of phagocytosis that is highly
	conserved across evolution and is mediated by the concerted action of several molecules.
	MEGF10 is an engulfment receptor protein that localizes to the plasma membrane in a
	punctuated pattern. The MEGF10 protein shares structural similarities with nematode
	engulfment receptor cell death abnormal-1 (CED-1). MEGF10 function is modulated by the ATP
	binding cassette transporter ABCA1, which is an ortholog to CED-7. It is expressed at the cell
	surface in clusters around cell corpses during engulfment, and during the engulfment of
	apoptotic thymocytes is recruited to the bottom of the forming phagocytic cup.
Gene ID:	84466
Pathways:	Regulation of Muscle Cell Differentiation
Application Details	
Application Notes:	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Torride.	
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish
	peroxidase.
Storage:	-20 °C

### Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months