# antibodies .- online.com





# anti-GLT25D2 antibody (AA 501-600) (HRP)



( )	1/0	r\ /1	014	
( )	ve	I V I	-v	V

Quantity:	100 μL	
Target:	GLT25D2	
Binding Specificity:	AA 501-600	
Reactivity:	Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GLT25D2 antibody is conjugated to HRP	
Application:	ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human GLT25D2	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Cow,Sheep,Pig,Horse,Chicken	
Purification:	Purified by Protein A.	

## Target Details

Target:	GLT25D2
Alternative Name:	GLT25D2 (GLT25D2 Products)

## Target Details

Background:	Synonyms: C1orf17, Glt25d2, Glycosyltransferase 25 family member 2, GT252_HUMAN,		
	Hydroxylysine galactosyltransferase 2, Procollagen galactosyltransferase 2, RP11-498P10.2.		
	Background: GLT25D2 is a 626 amino acid protein that localizes to endoplasmic reticulum		
	lumen and belongs to the glycosyltransferase 25 family. Possessing beta-galactosyltransferase		
	activity, GLT25D2 transports beta-galactose to hydroxylysine residues on collagen but lacks		
	glucosyltransferase activity. GLT25D2 is expressed in skeletal muscle and brain, and is		
	encoded by a gene located on human chromosome 1q25.3, a region associated with the		
	hereditary prostate cancer (HPC1) locus. Human chromosome 1 spans 260 million base pairs,		
	contains over 3,000 genes, comprises nearly 8 % of the human genome and houses a large		
	number of disease-associated genes, including those that are involved in familial adenomatous		
	polyposis, Stickler syndrome, Parkinson?s disease, Gaucher disease, schizophrenia and Usher		
	syndrome.		
Gene ID:	23127		
Application Details			
Application Notes:	IHC-P 1:200-400		
	IHC-F 1:100-500		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish		
	peroxidase.		
Storage:	-20 °C		
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.		

	lI	1:
-	เวทก	111111
	land	11110

Expiry Date:

12 months