



[Go to Product page](#)

Datasheet for ABIN1711904
anti-IB2 antibody (AA 31-130) (HRP)

Overview

Quantity:	100 µL
Target:	IB2 (JIP-2)
Binding Specificity:	AA 31-130
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IB2 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human JIP2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Horse
Purification:	Purified by Protein A.

Target Details

Target:	IB2 (JIP-2)
Alternative Name:	JIP2 (JIP-2 Products)
Background:	Synonyms: MAPK8IP2, C jun amino terminal kinase interacting protein 2, C-jun-amino-terminal

Target Details

kinase-interacting protein 2, Homologous to mouse JIP 1, IB 2, IB-2, IB2, Islet brain 2, Islet-brain-2, JIP 2, JIP-2, JIP2, JIP2_HUMAN, JNK interacting protein 2, JNK MAP kinase scaffold protein 2, JNK MAP kinase scaffold protein JIP2, JNK-interacting protein 2, MAPK8IP2, Mitogen activated protein kinase 8 interacting protein 2, Mitogen-activated protein kinase 8-interacting protein 2, PRKM8 interacting protein like, PRKM8IPL.

Background: c-Jun NH2-terminal kinases (JNKs) are distant members of the MAP kinase family (1). JNK1 is activated by dual phosphorylation at a Thr-Pro-Tyr motif in response to ultraviolet (UV) light, and it functions to phosphorylate c-Jun at amino terminal serine regulatory sites, Ser-63 and Ser-73, resulting in transcriptional activation (2-5). Two additional JNK family members have been identified as JNK2 and JNK3 (3). JIP-1 (for JNK interacting protein-1) has been identified as a cytoplasmic inhibitor of JNK that retains JNK in the cytoplasm, thereby inhibiting JNK-regulated gene expression. Evidence suggests that JNK1 and JNK2 bind to JIP-1 with greater affinity than to ATF-2 and c-Jun, which are targets of the JNK signaling pathway. JIP-1 contains an amino terminal JNK binding domain and a carboxy terminal SH3 domain. ATF-2 and c-Jun also contain the JNK binding domain and are thought to compete with JIP-1 for JNK binding (6). Multiple splice variants of JIP-1, including JIP-1b, JIP-1c (also designated islet-brain 1 or IB-1), JIP-2a, JIP-2b and JIP-3, have been identified in brain (7).

Application Details

Application Notes: WB 1:300-5000
 IHC-P 1:200-400
 IHC-F 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish

Handling

peroxidase.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months