# antibodies .- online.com







## anti-FANCE antibody (AA 431-536) (HRP)



| ( ) | 11/0   | K\ /   | iew         |
|-----|--------|--------|-------------|
|     | $\cup$ | 'I V/I | $I \cap VV$ |
|     |        |        |             |

| Quantity:            | 100 μL  |  |
|----------------------|---|--|
| Target:              | FANCE   |  |
| Binding Specificity: | AA 431-536  |  |
| Reactivity:          | Human   |  |
| Host:                | Rabbit  |  |
| Clonality:           | Polyclonal  |  |
| Conjugate:           | This FANCE antibody is conjugated to HRP  |  |
| Application:         | Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)) |  |

#### **Product Details**

| Immunogen:            | KLH conjugated synthetic peptide derived from human FANCE |
|-----------------------|---|
| Isotype:              | IgG   |
| Predicted Reactivity: | Human,Mouse,Rat,Dog                                       |
| Purification:         | Purified by Protein A.                                    |

#### **Target Details**

| Target:           | FANCE   |  |
|-------------------|---|--|
| Alternative Name: | FANCE (FANCE Products)  |  |
| Background:       | ackground: Synonyms: 2810451D06Rik, Al415634, AW209126, FACE, FAE, FANCE_HUMAN, |  |

Fanconi anemia complementation group E, Fanconi anemia group E protein, Protein FACE, RGD1561045.

Background: The Fanconi anemia complementation group (FANC) currently includes FANCA, FANCB, FANCC, FANCD1 (also called BRCA2), FANCD2, FANCE, FANCF, FANCG, FANCI, FANCJ (also called BRIP1), FANCL, FANCM and FANCN (also called PALB2). The previously defined group FANCH is the same as FANCA. Fanconi anemia is a genetically heterogeneous recessive disorder characterized by cytogenetic instability, hypersensitivity to DNA crosslinking agents, increased chromosomal breakage, and defective DNA repair. The members of the Fanconi anemia complementation group do not share sequence similarity, they are related by their assembly into a common nuclear protein complex. This gene encodes the protein for complementation group E. [provided by RefSeq, Jul 2008].

Gene ID: 2178

Pathways: DNA Damage Repair

#### **Application Details**

Application Notes: WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions: For Research Use only

#### Handling

| Format:            | Liquid   |
|--------------------|--|
| Concentration:     | 1 μg/μL  |
| Buffer:            | Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.         |
| Preservative:      | ProClin  |
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Handling Advice:   | Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.               |
| Storage:           | -20 °C   |
|                    |  |

### Handling

| Storage Comment: | Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles. |
|------------------|---|
| Expiry Date:     | 12 months   |