antibodies -online.com





Datasheet for ABIN1712130 anti-FOXN2 antibody (AA 51-150) (HRP)

Go to Product page

Overview	
Quantity:	100 μL
Target:	FOXN2
Binding Specificity:	AA 51-150
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FOXN2 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human FOXN2
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Pig,Rabbit
Purification:	Purified by Protein A.
Target Details	
Target:	FOXN2
Alternative Name:	FOXN2 (FOXN2 Products)

Target Details

Background:

Synonyms: 3230402J05Rik, 6030465J18Rik, Fkh19, Forkhead box N2, Forkhead box protein N2, Foxn1, foxn2, FOXN2_HUMAN, HTLF, Human T-cell leukemia virus enhancer factor. Background: The forkhead domain-containing gene family (Fox) comprises over 20 members in mammals and is defined by a conserved 110 amino-acid motif containing a winged helix structure DNA-binding domain. The members of this gene family have been implicated as key regulators of embryogenesis, cell cycling, cell lineage restriction and cancer. As such, FOXN2 contains a domain with homology to the forkhead DNA binding domain. FOXN2, or Human T-cell leukemia virus enhancer factor, is a 341 amino acid protein mapping to human gene FOXN2, which has been localized to human chromosome 2p16-p22. This protein, encoded by a 1239-bp cDNA isolated from the Jurkat cDNA library, is capable of binding to a region of the human T-cell leukemia virus long terminal repeat (HTLV-I LTR) located between amino acids 155 and 117. This purine-rich region is important in the regulation of gene expression by the Ets family of transcription factors. FOXN2 is a unique cellular gene that may function in the transcriptional regulation of HTLV-I LTR.

Gene ID:

3344

Application Details

Λnn	lication	Motae.
\neg	IICALIOLE	TAUTES.

WB 1:300-5000

IHC-P 1:200-400

IHC-F 1:100-500

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months