

## Datasheet for ABIN1712332

## anti-GNPDA2 antibody (AA 101-200) (HRP)



Go to Product page

_					
	W	0	rv	10	W

Quantity:	100 μL	
Target:	GNPDA2	
Binding Specificity:	AA 101-200	
Reactivity:	Mouse, Rat	
Host:	Rabbit	
Clonality:	Polyclonal	
Conjugate:	This GNPDA2 antibody is conjugated to HRP	
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))	
Product Details		
Immunogen:	KLH conjugated synthetic peptide derived from human GNPDA2	
Isotype:	IgG	
Cross-Reactivity:	Mouse, Rat	
Predicted Reactivity:	Human,Dog,Cow,Sheep,Pig,Horse	
Purification:	Purified by Protein A.	
Target Details		
Target:	GNPDA2	
Alternative Name:	GNPDA2 (GNPDA2 Products)	

## **Target Details**

Background:	Synonyms: GlcN6P deaminase 2, Glucosamine 6 phosphate deaminase 2, Glucosamine 6		
	phosphate isomerase 2, Glucosamine 6 phosphate isomerase SB52, Glucosamine-6-phosphate		
	deaminase 2, Glucosamine-6-phosphate isomerase 2, Glucosamine-6-phosphate isomerase		
	SB52, GNP2, GNPDA 2, Gnpda2, GNPI2_HUMAN, SB52.		
	Background: During fertilization in mammals, the sperm activates the egg by causing an		
	increase in the level of free cytoplasmic calcium concentration. This increased calcium		
	concentration induces a characteristic series of oscillations that trigger egg activation and early		
	embryo development. A hamster protein named oscillin is thought to be involved in this		
	pathway. The enzyme glucosamine-6-phosphate isomerase (GNPI) or deaminase (GNPDA1)		
	and the related protein GNPDA2 are the human homologs of hamster oscillin. GNPDA1 and		
	GNPDA2 catalyze the conversion of GNP to fructose-6-phosphate and ammonia. Both proteins		
	exist as homohexamers and are ubiquitously expressed with highest expression in testis, ovary		
	and heart. Three isoforms of GNPDA2 are expressed due to alternative splicing events.		
Gene ID:	132789		
Application Dataila			
Application Details			
Application Notes:	WB 1:300-5000		
	IHC-P 1:200-400		
	IHC-F 1:100-500		
Restrictions:	For Research Use only		
Handling			
Format:	Liquid		
Concentration:	1 μg/μL		
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and		
	50 % Glycerol.		
Preservative:	ProClin		
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be		
	handled by trained staff only.		
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish		
	peroxidase.		
Storage:	-20 °C		

## Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months