

Datasheet for ABIN1712412

## anti-ZBTB4 antibody (AA 301-400) (HRP)



[Go to Product page](#)

### Overview

|                      |  |
|----------------------|--|
| Quantity:            | 100 µL   |
| Target:              | ZBTB4  |
| Binding Specificity: | AA 301-400   |
| Reactivity:          | Human  |
| Host:                | Rabbit   |
| Clonality:           | Polyclonal   |
| Conjugate:           | This ZBTB4 antibody is conjugated to HRP   |
| Application:         | Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)),<br>Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)) |

### Product Details

|                       |  |
|-----------------------|--|
| Immunogen:            | KLH conjugated synthetic peptide derived from human ZBTB4/ZNF903 |
| Isotype:              | IgG  |
| Predicted Reactivity: | Human, Mouse, Rat, Dog, Cow, Sheep, Pig, Horse, Rabbit           |
| Purification:         | Purified by Protein A.   |

### Target Details

|                   |  |
|-------------------|--|
| Target:           | ZBTB4  |
| Alternative Name: | ZBTB4/ZNF903 ( <a href="#">ZBTB4 Products</a> )  |
| Background:       | Synonyms: KAISO-L1, KAISO-like zinc finger protein 1, KIAA1538, ZBTB4, ZBTB4_HUMAN, Zinc |

## Target Details

finger and BTB domain containing 4, Zinc finger and BTB domain-containing protein 4, ZNF903.

Background: Zinc-finger proteins contain DNA-binding domains and have a wide variety of functions, most of which encompass some form of transcriptional activation or repression. The majority of zinc-finger proteins contain a Kr<sub>ppel</sub>-type DNA binding domain and a KRAB domain, which is thought to interact with KAP1, thereby recruiting histone modifying proteins.

ZBTB4 (zinc finger and BTB domain containing 4), also known as KAISO-L1 (KAISO-like zinc finger protein 1), is a 1,013 amino acid nuclear protein that is involved in transcriptional regulation. ZBTB4 contains one BTB (POZ) domain, six C2H2-type zinc fingers and is phosphorylated and downregulated by HIPK2. The gene encoding ZBTB4 maps to human chromosome 17, which comprises over 2.5 % of the human genome and encodes over 1,200 genes.

Gene ID: 57659

## Application Details

Application Notes: WB 1:300-5000  
IHC-P 1:200-400  
IHC-F 1:100-500

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: Aqueous buffered solution containing 0.01M TBS ( pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Handling Advice: Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish peroxidase.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

## Handling

---

Expiry Date: 12 months