

### Datasheet for ABIN1712545

# anti-GOLGA5 antibody (AA 401-500) (HRP)

GOLGA5

GOLGA5 (GOLGA5 Products)



#### Overview

Target:

Alternative Name:

Overview	
Quantity:	100 μL
Target:	GOLGA5
Binding Specificity:	AA 401-500
Reactivity:	Mouse, Rat
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This GOLGA5 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))
Product Details	
Immunogen:	KLH conjugated synthetic peptide derived from human GOLGA5
Isotype:	IgG
Cross-Reactivity:	Mouse, Rat
Predicted Reactivity:	Human,Cow,Pig,Horse,Rabbit
Purification:	Purified by Protein A.
Target Details	

### Target Details

Background:	Synonyms: Cell proliferation inducing gene 31, Cell proliferation-inducing gene 31 protein,	
	GOGA5_HUMAN, GOLGA 5, GOLGA5, Golgi autoantigen golgin subfamily a 5, Golgi integral	
	membrane protein 5, Golgin 84, Golgin subfamily A member 5, Golgin-84, Golgin84, GOLIM 5,	
	GOLIM5, Protein Ret-II, PTC5, Ret fused gene 5, RET fused gene 5 protein, Ret II, Ret II protein,	
	RET-fused gene 5 protein, RFG 5, RFG5.	
	Background: The Golgi apparatus, which participates in glycosylation and transport of proteins	
	and lipids in the secretory pathway, consists of a series of stacked cisternae (flattened	
	membrane sacs). Interactions between the Golgi and microtubules are thought to be important for the reorganization of the Golgi after it fragments during mitosis. This gene encodes one of the golgins, a family of proteins localized to the Golgi. This protein is a coiled-coil membrane	
		protein that has been postulated to play a role in vesicle tethering and docking. Translocations
		involving this gene and the ret proto-oncogene have been found in tumor tissues, the chimeric
	sequences have been designated RET-II and PTC5. [provided by RefSeq, Feb 2010].	
	Gene ID:	9950
Application Details		
Application Notes:	WB 1:300-5000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and	
	50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be	
	handled by trained staff only.	
Handling Advice:	Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish	
	peroxidase.	
Storage:	-20 °C	

# Handling

Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months