

## Datasheet for ABIN1712560 anti-MIXL1 antibody (AA 101-200) (HRP)



Go to Product page

Overview	
Quantity:	100 μL
Target:	MIXL1
Binding Specificity:	AA 101-200
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This MIXL1 antibody is conjugated to HRP
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)),
	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p))
Product Details	
Product Details Immunogen:	KLH conjugated synthetic peptide derived from human MIXL1
	KLH conjugated synthetic peptide derived from human MIXL1
Immunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Cross-Reactivity:	IgG Mouse
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity:	IgG  Mouse  Human,Rat,Cow,Sheep,Pig,Horse
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity: Purification:	IgG  Mouse  Human,Rat,Cow,Sheep,Pig,Horse

## **Target Details**

Handling Advice:

Target Details	
Background:	Synonyms: hMix, Homeobox protein MIXL1, Homeodomain protein 1, Homeodomain protein
	MIX, MILD1, MIX, Mix.1, Mix.1 homeobox-like protein, Mix1 homeobox-like 1, MIX1 homeobox-
	like protein 1, MixI1, MIXL1_HUMAN.
	Background: The homeobox DNA-binding domain is a 60 amino acid motif that is conserved
	among many species and functions to bind DNA via a helix-turn-helix structure, thereby playing
	a role in transcriptional regulation and in the control of gene expression. MIXL1 (Mix1
	homeobox-like 1), also known as MIXL, is a 232 amino acid protein that localizes to the nucleus
	and contains one homeobox DNA-binding domain. Expressed in lymph tissues, MIXL1
	functions as a transcription factor that plays an essential role in axial mesendoderm
	morphogenesis and endoderm formation and is also required for cellular differentiation during
	blood development. Additionally, MIXL1 is involved in maturation of heart and gut tissue during
	embryogenesis and may also act as a negative regulator of brachyury expression.
	Overexpression of MIXL1 is associated with non-Hodgkin and Hodgkin lymphomas, suggesting
	a role in carcinogenesis.
Gene ID:	83881
Application Details	
Application Notes:	WB 1:300-5000
	IHC-P 1:200-400
	IHC-F 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and
	50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be
	handled by trained staff only.

peroxidase.

Do NOT add Sodium Azide! Use of Sodium Azide will inhibit enzyme activity of horseradish

## Handling

Storage:	-20 °C
Storage Comment:	Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.
Expiry Date:	12 months