

Datasheet for ABIN1713153 **anti-HFE antibody (PE-Cy5.5)**



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	HFE
Reactivity:	Human, Mouse, Rat, Cow, Dog
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This HFE antibody is conjugated to PE-Cy5.5
Application:	Western Blotting (WB), Flow Cytometry (FACS)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human HFE/Hemochromatosis
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Dog,Cow,Sheep,Horse
Purification:	Purified by Protein A.

Target Details

Target:	HFE
Alternative Name:	HFE/Hemochromatosis (HFE Products)
Background:	Synonyms: dJ221C16.10.1, Hemochromatosis, Hemochromatosis protein, Hereditary hemochromatosis protein, Hereditary hemochromatosis protein HLA H, HFE 1, HFE,

Target Details

HFE_HUMAN, HFE1, HH, High Fe, HLA H, HLA-H, HLAH, MGC:150812, MGC10379, MGC103790, MHC class I like protein HFE, MVCD7, TFQTL2.

Background: The features of hemochromatosis include cirrhosis of the liver, diabetes, hypermelanotic pigmentation of the skin, and heart failure. Since hemochromatosis is a relatively easily treated disorder if diagnosed, this is a form of preventable cancer. The HFE protein, which is defective in hereditary hemo-chromatosis, normally is expressed in crypt enterocytes of the duodenum where it has a unique, predominantly intracellular localization. In placenta, the HFE protein co-localizes with and forms a stable association with the transferrin receptor (TfR), providing a link between the HFE protein and iron transport.

Immunocytochemistry shows that the HFE protein and TfR both are expressed in the crypt enterocytes. Western blots show that, as is the case in human placenta, the HFE protein in crypt enterocytes is physically associated with the TfR and with α_2 -microglobulin. It is proposed that HFE has two mutually exclusive activities in cells: inhibition of uptake or inhibition of release of iron and that the balance between serum transferrin saturation and serum transferrin-receptor concentrations determines which of these functions predominates. The gene which encodes HFE maps to human chromosome 6p21.3.

Gene ID: 3077

Pathways: [Transition Metal Ion Homeostasis](#), [Regulation of Leukocyte Mediated Immunity](#), [Positive Regulation of Immune Effector Process](#)

Application Details

Application Notes: FCM(1:100-500)
Optimal working dilution should be determined by the investigator.

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 $\mu\text{g}/\mu\text{L}$

Buffer: Aqueous buffered solution containing 0.01M TBS (pH 7.4) with 1 % BSA, 0.03 % Proclin300 and 50 % Glycerol.

Preservative: Sodium azide

Precaution of Use: This product contains sodium azide: a POISONOUS AND HAZARDOUS SUBSTANCE which

Handling

should be handled by trained staff only.

Storage: -20 °C

Storage Comment: Store at -20°C. Aliquot into multiple vials to avoid repeated freeze-thaw cycles.

Expiry Date: 12 months