# antibodies -online.com





## anti-TIFA antibody (AA 51-150)



Go to Product page

$\sim$					
()	VE	۲۱	/1	$\triangle$	Λ

Quantity:	100 μL
Target:	TIFA
Binding Specificity:	AA 51-150
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TIFA antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC)

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human TIFA	
Isotype:	IgG	
Cross-Reactivity:	Mouse	
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Chicken	
Purification:	Purified by Protein A.	

#### **Target Details**

Target: TIFA	
--------------	--

### **Target Details**

Expiry Date:

Alternative Name:	TIFA (TIFA Products)	
Background:	Synonyms: Putative MAPK activating protein PM14, Putative MAPK-activating protein PM14, Putative NF kappa B activating protein 20, Putative NF-kappa-B-activating protein 20, T2BP, TA, TA_HUMAN, TRAF interacting protein with FHA domain containing protein A, TRAF-interacting protein with FHA domain-containing protein A, TRAF2 binding protein, TRAF2-binding protein. Background: Adapter protein which mediates the IRAK1 and TRAF6 interaction following IL-1 stimulation, resulting in the downstream activation of NF-kappa-B and AP-1 pathways. Induces the oligomerization and polyubiquitination of TRAF6, which leads to the activation of TAK1 and IKK through a proteasome-independent mechanism.	
Application Details		
Application Notes:  Restrictions:	WB 1:300-5000  ELISA 1:500-1000  IHC-P 1:200-400  IHC-F 1:100-500  IF(IHC-P) 1:50-200  IF(IHC-F) 1:50-200  IF(ICC) 1:50-200  ICC 1:100-500  For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	

12 months