



[Go to Product page](#)

Datasheet for ABIN1713659
anti-IdnK antibody (AA 51-150)

Overview

| | |
|----------------------|---|
| Quantity: | 100 µL |
| Target: | IdnK (IDNK) |
| Binding Specificity: | AA 51-150 |
| Reactivity: | Human |
| Host: | Rabbit |
| Clonality: | Polyclonal |
| Conjugate: | This IdnK antibody is un-conjugated |
| Application: | Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro)) |

Product Details

| | |
|-----------------------|--|
| Immunogen: | KLH conjugated synthetic peptide derived from human C9orf103 |
| Isotype: | IgG |
| Predicted Reactivity: | Human,Mouse |
| Purification: | Purified by Protein A. |

Target Details

| | |
|-------------------|--|
| Target: | IdnK (IDNK) |
| Alternative Name: | C9orf103 (IDNK Products) |

Target Details

Background: Synonyms: bA522I20.2, C9orf103, Chromosome 9 open reading frame 103, Glucokinase like protein, Gluconate kinase, Gluconokinase like protein, GNTK_HUMAN, IDNK, OTTHUMP00000021546, OTTHUMP00000021547, Probable gluconokinase, RP11-522I20.2. Background: C9orf103 (chromosome 9 open reading frame 103), also known as gluconate kinase, is a 187 amino acid protein that belongs to the gluconokinase gntK/gntV family and catalyzes the conversion of ATP and D-gluconate to ADP and 6-D-gluconate. Existing as three alternatively spliced isoforms, the gene encoding C9orf103 maps to human chromosome 9q21.32. Chromosome 9 consists of about 145 million bases, represents 4 % of the human genome and encodes nearly 900 genes. Thought to play a role in gender determination, deletion of the distal portion of 9p can lead to development of male to female sex reversal, the phenotype of a female with a male X,Y genotype. Hereditary hemorrhagic telangiectasia, which is characterized by harmful vascular defects, is associated with the chromosome 9 gene encoding endoglin protein, ENG. Familial dysautonomia is also associated with chromosome 9 though through the gene IKBKAP. Notably, chromosome 9 encompasses the largest interferon family gene cluster.

Gene ID: 414328

Application Details

Application Notes: WB 1:300-5000
ELISA 1:500-1000
IHC-P 1:200-400
IHC-F 1:100-500
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
ICC 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Handling

| | |
|--------------------|--|
| Precaution of Use: | This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only. |
| Storage: | 4 °C,-20 °C |
| Storage Comment: | Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles. |
| Expiry Date: | 12 months |