

Datasheet for ABIN1713852

anti-IRF6 antibody (AA 1-100)



Overview

Quantity:	100 μL
Target:	IRF6
Binding Specificity:	AA 1-100
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This IRF6 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Dot Blot (DB)
Product Details	
1 Toddot Details	
Immunogen:	KLH conjugated synthetic peptide derived from human IRF6
	KLH conjugated synthetic peptide derived from human IRF6
Immunogen:	
Immunogen: Isotype:	IgG
Immunogen: Isotype: Cross-Reactivity:	IgG Human, Mouse
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity:	IgG Human, Mouse Rat,Cow,Sheep,Pig,Rabbit
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity: Purification:	IgG Human, Mouse Rat,Cow,Sheep,Pig,Rabbit
Immunogen: Isotype: Cross-Reactivity: Predicted Reactivity: Purification: Target Details	IgG Human, Mouse Rat,Cow,Sheep,Pig,Rabbit Purified by Protein A.

OTTHUMP00000034677, OTTHUMP00000034678, PIT, PPS, PPS1, VWS, VWS1.

Background: Interferon regulatory factor-1 (IRF-1) and IRF-2 have been identified as novel DNA-binding factors that function as regulators of both type I interferon (interferon-alpha and beta) and interferon-inducible genes. The two factors are structurally related, particularly in their N-terminal regions, which confer DNA binding specificity. In addition, both bind to the same sequence within the promoters of interferon-alpha and interferon-beta genes. IRF-1 functions as an activator of interferon transcription, while IRF-2 binds to the same cis elements and represses IRF-1 action. IRF-1 and IRF-2 have been reported to act in a mutually antagonistic manner in regulating cell growth, overexpression of the repressor IRF-2 leads to cell transformation while concomitant overexpression of IRF-1 causes reversion. IRF-1 and IRF-2 are members of a larger family of DNA binding proteins that includes IRF-3, IRF-4, IRF-5, IRF-6, IRF-7, ISGF-3 gamma p48 and IFN consensus sequence-binding protein (ICSBP).

Application Details

Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	dot-blot()
	V

Restrictions: For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months