



[Go to Product page](#)

Datasheet for ABIN1713868
anti-NCKAP5 antibody (AA 1-100)

Overview

Quantity:	100 µL
Target:	NCKAP5
Binding Specificity:	AA 1-100
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This NCKAP5 antibody is un-conjugated
Application:	Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human NCKAP5
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Horse,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	NCKAP5
Alternative Name:	NCKAP5 (NCKAP5 Products)

Target Details

Background:	<p>Synonyms: ERIH1, ERIH2, NAP-5, NAP5, Nck-associated protein 5, NCKAP5, NCKP5_HUMAN, Peripheral clock protein.</p> <p>Background: NAP5 (Nck-associated protein 5), also known as peripheral clock protein, NCKAP5 or ERIH, is a 1,909 amino acid nuclear protein that is expressed in fetal and adult brain, leukocytes and fetal fibroblasts. Containing pro-rich sequences, NAP5 interacts with the adapter protein Nck via the SH3-containing region. Existing as four alternatively spliced isoforms, the gene encoding NAP5 maps to human chromosome 2q21.2 and mouse chromosome 1 E3. Human chromosome 2, the second largest human chromosome, consists of 237 million bases, encodes over 1,400 genes and makes up approximately 8 % of the human genome. Harlequin ichthyosis, a rare and morbid skin deformity, is associated with mutations in the ABCA12 gene present on chromosome 2. The lipid metabolic disorder sitosterolemia is associated with ABCG5 and ABCG8. An extremely rare recessive genetic disorder, Alstr syndrome, is due to mutations in the ALMS1 gene.</p>
-------------	--

Application Details

Application Notes:	<p>ELISA 1:500-1000</p> <p>IHC-P 1:200-400</p> <p>IHC-F 1:100-500</p> <p>IF(IHC-P) 1:50-200</p> <p>IF(IHC-F) 1:50-200</p> <p>IF(ICC) 1:50-200</p> <p>ICC 1:100-500</p>
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C

Handling

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date: 12 months