

[Go to Product page](#)

## Datasheet for ABIN1713885 **anti-PRELP antibody (AA 151-250)**

### Overview

Quantity:	100 µL
Target:	PRELP
Binding Specificity:	AA 151-250
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This PRELP antibody is un-conjugated
Application:	Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human PRELP
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Horse,Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	PRELP
Alternative Name:	PRELP ( <a href="#">PRELP Products</a> )
Background:	Synonyms: 55 kDa leucine rich repeat protein of articular cartilage, MST161, MSTP161, Prelp,

## Target Details

PRELP\_HUMAN, Prolargin, Prolargin proteoglycan, Proline arginine rich end leucine rich repeat protein, Proline-arginine-rich end leucine-rich repeat protein, Proline/arginine rich end leucine rich repeat protein, SLRR2A.

Background: PRELP (proline/arginine-rich end leucine-rich repeat protein), also known as Prolargin, MST161, SLRR2A or MSTP161, is a 382 amino acid secreted protein that localizes to the extracellular matrix. Belonging to the Class II subfamily of the small leucine-rich proteoglycan (SLRP) family, PRELP contains twelve LRR (leucine-rich) repeats, which are motifs consisting of 20-29 residues that are present in numerous proteins with diverse functions and provide versatile structural framework for the formation of protein-protein interactions. Highly expressed in cartilage, basement membranes and developing bone, PRELP is considered a glycosaminoglycan (GAG)- and collagen-binding anchor protein that associates with the basement membrane heparan sulfate proteoglycan perlecan. PRELP acts as a linker between the extracellular matrix and the cell surface of proteoglycans and may be partially responsible for Hutchinson-Gilford progeria (HGP), an extremely rare genetic disorder that causes premature, rapid aging shortly after birth.

Pathways: [Glycosaminoglycan Metabolic Process](#)

## Application Details

Application Notes: ELISA 1:500-1000

Restrictions: For Research Use only

## Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Preservative: ProClin

Precaution of Use: This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.

Storage: 4 °C, -20 °C

Storage Comment: Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.

Expiry Date: 12 months