antibodies -online.com





TETIA KENINEWEA

Datasheet for ABIN1713916

anti-ANTXR1 antibody (AA 201-300)

Overview

Quantity:	100 μL
Target:	ANTXR1
Binding Specificity:	AA 201-300
Reactivity:	Human, Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ANTXR1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TEM8
Isotype:	IgG
Cross-Reactivity:	Human, Mouse
Predicted Reactivity:	Rat,Rabbit
Purification:	Purified by Protein A.

Target Details

Target: ANTXR1

Target Details

Alternative Name:	TEM8 (ANTXR1 Products)
Background:	Synonyms: Anthrax toxin receptor 1, Anthrax toxin receptor 1 precursor, ANTXR 1, ANTXR1,
	ATR, TEM-8, TEM 8, TEM8, Tumor Endothelial Marker 8, Tumor endothelial marker 8 precursor,
	ANTR1_HUMAN, Anthrax toxin receptor-1, FLJ11298, FLJ21776.
	Background: This gene encodes a type I transmembrane protein and is a tumor-specific
	endothelial marker that has been implicated in colorectal cancer. The encoded protein has beer
	shown to also be a docking protein or receptor for Bacillus anthracis toxin, the causative agent
	of the disease, anthrax. The binding of the protective antigen (PA) component, of the tripartite
	anthrax toxin, to this receptor protein mediates delivery of toxin components to the cytosol of
	cells. Once inside the cell, the other two components of anthrax toxin, edema factor (EF) and
	lethal factor (LF) disrupt normal cellular processes. Three alternatively spliced variants that
	encode different protein isoforms have been described. [provided by RefSeq, Oct 2008]
Gene ID:	84168
UniProt:	Q9H6X2
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin

Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months