antibodies - online.com







anti-FAF2 antibody (AA 81-180)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	FAF2
Binding Specificity:	AA 81-180
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FAF2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ETEA
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Sheep,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	FAF2		
---------	------	--	--

Target Details

Alternative Name:	ETEA (FAF2 Products)
Background:	Synonyms: Al462440, expressed in T cells and eosinophils in atopic dermatitis, faf2,
	FAF2_HUMAN, Fas associated factor family member 2, FAS-associated factor 2, KIAA0887,
	Protein ETEA, Protein expressed in T cells and eosinophils in atopic dermatitis, Protein
	expressed in T-cells and eosinophils in atopic dermatitis, UBX domain containing 8, UBX
	domain protein 3B, UBX domain-containing protein 3B, UBX domain-containing protein 8, UBXD8, UBXN3B.
	Background: ETEA is a 445-amino acid protein associated with atopic dermatitis (AD), a chronic
	noncontagious relapsing inflammatory skin disease characterized by eczematous skin lesions
	and also referred to as eczematous dermatitis. Other atopic diseases such as hay fever,
	asthma and conjunctivitis often occur along with AD. ETEA shows higher expression in T cells
	and eosinophils of patients with AD than in T cells and eosinophils of unaffected individuals. T
	cells are influential in the regulation of the inflammatory process of this disease. The
	persistence of jAD is attributed to dysregulated apoptosis in T cells, eosinophils, and
	keratinocytes. ETEA may be involved in the resistance to apoptosis in T cells and eosinophils o
	AD patients.
Gene ID:	23197
Application Details	
	WB 1:300-5000
	WB 1:300-5000 ELISA 1:500-1000
	ELISA 1:500-1000
	ELISA 1:500-1000 IHC-P 1:200-400
	ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500
	ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200
	ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200
Application Notes:	ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200
Application Notes: Restrictions:	ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
Application Details Application Notes: Restrictions: Handling Format:	ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
Application Notes: Restrictions: Handling	ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500 For Research Use only

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months