

[Go to Product page](#)

Datasheet for ABIN1713977 **anti-TESK2 antibody (AA 301-400)**

Overview

Quantity:	100 µL
Target:	TESK2
Binding Specificity:	AA 301-400
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TSK2 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin- embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human TSK2
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Horse,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	TESK2
Alternative Name:	TESK2 (TESK2 Products)

Target Details

Background:	<p>Synonyms: Dual specificity testis specific protein kinase 2, Dual specificity testis-specific protein kinase 2, MGC29168, OTTHUMP00000009093, RP23-109A3.2, Tesk2, TESK2_HUMAN, Testicular protein kinase 2, Testis specific kinase 2.</p> <p>Background: TESK2 is a nuclear protein that belongs to the protein kinase superfamily and is expressed in testis and prostate. Functioning as a dual-specificity protein kinase, TESK2 catalyzes the ATP-dependent phosphorylation of substrates and autophosphorylation on tyrosine and serine/threonine residues, thereby mediating intracellular signal transduction pathways. TESK2 requires magnesium as a cofactor and its catalytic activity is thought to play an important role in meiotic events such as spermatogenesis. TESK2 contains one protein kinase domain that is 65 % identical to the kinase domain found in TESK1 (testicular protein kinase 1), suggesting a similar role for these proteins in phosphorylation events. Three isoforms of TESK2 are expressed due to alternative splicing.</p>
Gene ID:	10420
Pathways:	Cell-Cell Junction Organization

Application Details

Application Notes:	<p>WB 1:300-5000</p> <p>ELISA 1:500-1000</p> <p>IHC-P 1:200-400</p> <p>IHC-F 1:100-500</p> <p>IF(IHC-P) 1:50-200</p> <p>IF(IHC-F) 1:50-200</p> <p>IF(ICC) 1:50-200</p> <p>ICC 1:100-500</p>
Restrictions:	For Research Use only

Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months