
antibodies - online.com







anti-C3orf58 antibody (AA 351-430)



Image



Overview

Quantity:	100 μL
Target:	C3orf58 (C3ORF58)
Binding Specificity:	AA 351-430
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This C3orf58 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human C3orf58
Isotype:	IgG
Cross-Reactivity:	Human
Predicted Reactivity:	Mouse,Rat,Cow,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	C3orf58 (C3ORF58)	
---------	-------------------	--

Target Details

Alternative Name:	C3orf58 (C3ORF58 Products)
Background:	Synonyms: Chromosome 3 open reading frame 58, DIA1, Golgi protein GoPro49, MGC33365,
	UPF0672 protein C3orf58, DIA1_HUMAN.
	Background: C3orf58 (chromosome 3 open reading frame 58), also known as DIA1, is a 430
	amino acid secreted protein that belongs to the UPF0672 family. C3orf58 is encoded by a gene
	that maps to human chromosome 3q24. Chromosome 3 is made up of approximately 214
	million bases encoding over 1,100 genes. Notably, there is a chemokine receptor gene cluster
	and a variety of human cancer related loci on chromosome 3. Particular regions of the
	chromosome 3 short arm are deleted in many types of cancer cells. Key tumor suppressing
	genes on chromosome 3 encode apoptosis mediator RASSF1, cell migration regulator HYAL1
	and angiogenesis suppressor SEMA3B. Marfan Syndrome, porphyria, von Hippel-Lindau
	syndrome, osteogenesis imperfecta and Charcot-Marie-Tooth disease are a few of the
	numerous genetic diseases associated with chromosome 3.
Gene ID:	205428
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months

Images




Image 1. Human TT lysates probed with C3orf58 Polyclonal Antibody, Unconjugated at 1:300 dilution and 4°C overnight incubation. Followed by conjugated secondary antibody incubation at 1:10000 for 60 min at 37°C.