# antibodies -online.com







# anti-ARID1B antibody (AA 1501-1600)



Image



#### Overview

Quantity:	100 μL
Target:	ARID1B
Binding Specificity:	AA 1501-1600
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ARID1B antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human ARID1B
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Chicken
Purification:	Purified by Protein A.

#### **Target Details**

et:	ARID1B		
-----	--------	--	--


## **Target Details**

Alternative Name:	ARID1B (ARID1B Products)
Background:	Synonyms: OSA2, 6A3-5, DAN15, MRD12, P250R, BRIGHT, BAF250B, ELD/OSA1, AT-rich
buonground.	interactive domain-containing protein 1B, ARID domain-containing protein 1B, BRG1-associated
	factor 250b, BRG1-binding protein hELD/OSA1, Osa homolog 2, hOsa2, ARID1B, KIAA1235
	Background: Involved in transcriptional activation and repression of select genes by chromatin
	remodeling (alteration of DNA-nucleosome topology). Belongs to the neural progenitors-
	specific chromatin remodeling complex (npBAF complex) and the neuron-specific chromatin
	remodeling complex (nBAF complex). During neural development a switch from a
	stem/progenitor to a post-mitotic chromatin remodeling mechanism occurs as neurons exit th
	cell cycle and become committed to their adult state. The transition from proliferating neural
	stem/progenitor cells to post-mitotic neurons requires a switch in subunit composition of the
	npBAF and nBAF complexes. As neural progenitors exit mitosis and differentiate into neurons,
	npBAF complexes which contain ACTL6A/BAF53A and PHF10/BAF45A, are exchanged for
	homologous alternative ACTL6B/BAF53B and DPF1/BAF45B or DPF3/BAF45C subunits in
	neuron-specific complexes (nBAF). The npBAF complex is essential for the self-
	renewal/proliferative capacity of the multipotent neural stem cells. The nBAF complex along
	with CREST plays a role regulating the activity of genes essential for dendrite growth (By
	similarity). Binds DNA non-specifically.
Gene ID:	57492
UniProt:	Q8NFD5
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only

### Handling

Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	

#### **Images**



#### **Immunohistochemistry (Paraffin-embedded Sections)**

**Image 1.** Formalin-fixed and paraffin embedded mouse embryo labeled with Rabbit Anti-ARID1B Polyclonal Antibody, Unconjugated (ABIN1714038) at 1:200 followed by conjugation to the secondary antibody and DAB staining