antibodies .- online.com







anti-ARFGEF1 antibody (AA 1-200)



()	11/0	K\ /	iew	1
	\cup	'I V/I	$\square \vee \vee$	ı

Quantity:	100 μL
Target:	ARFGEF1
Binding Specificity:	AA 1-200
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ARFGEF1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), ELISA, Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human BIG1/ARFGEF1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Chicken
Purification:	Purified by Protein A.

Target Details

Target:	ARFGEF1		
---------	---------	--	--

Target Details

Alternative Name:	BIG1/ARFGEF1 (ARFGEF1 Products)	
Background:	Synonyms: ADPribosylation factor guanine nucleotide exchange factor 1brefeldin A inhibited,	
	ARFGEF1, ARFGEP1, Brefeldin A inhibited GEP 1, Brefeldin A inhibited guanine nucleotide	
	exchange protein 1, p200 ARF GEP1, p200 ARF guanine nucleotide exchange factor, BIG1_HUMAN.	
	Background: Guanine nucleotide-exchange proteins (GEPs) accelerate replacement of bound	
	GDP with GTP and thereby activate ADP-ribosylation factors (ARFs), a family of guanine	
	nucleotide-binding proteins that play an important role in intracellular vesicular trafficking. GEPs	
	comprise two major families, large GEPs that are inhibited by brefeldin A (BFA), a protein that	
	effects Golgi structure and a group of smaller GEPs that are insenstive to BFA. Two genes for	
	GEPs found on human chromosomes 8 and 20 encode BFA sensitive GEPs designated BIG1	
	and BIG2. Both GEPS contain a sec7 domain that is responsible for their brefeldin inhibition and	
	also their catalytic activity. In vivo, BIG1 and BIG2 exist in macromolecular complexes that	
	move between the Golgi membranes and cytosol. BIG2 associates with PKA regulatory	
	subunits, implying that BIG2 may act as an A kinase-anchoring protein (AKAP) that could	
	coordinate the cAMP and ARF regulatory pathways.	
Gene ID:	10565	
Pathways:	Regulation of Actin Filament Polymerization, Regulation of Carbohydrate Metabolic Process	
Application Details		
Application Notes:	WB 1:300-5000	
	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	

Handling

Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	