# antibodies -online.com





Datasheet for ABIN1714117

## anti-BOP1 antibody (AA 201-300)



#### Overview

Quantity:	100 μL
Target:	BOP1
Binding Specificity:	AA 201-300
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This BOP1 antibody is un-conjugated
Application:	Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), ELISA, Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

#### **Product Details**

Immunogen:	KLH conjugated synthetic peptide derived from human BOP1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Chicken
Purification:	Purified by Protein A.

#### **Target Details**

Target:	BOP1		
---------	------	--	--

### **Target Details**

Alternative Name:	BOP1 (BOP1 Products)
Background:	Synonyms: AU020183, AW146150, Block of proliferation 1, Block of proliferation 1 protein,
	Bop1, Bop1 block of proliferation 1, BOP1_HUMAN, D18861, Erb1p, Kiaa0124, MGC109114,
	mKIAA0124, Ribosome biogenesis protein BOP1.
	Background: Predominantly localized to the nucleolus, BOP1 (Block of proliferation 1 protein) is
	a 746 amino acid highly conserved non-ribosomal protein that is involved in ribosome
	biogenesis. Truncation of the amino terminus of BOP1 leads to cell growth arrest in the G1
	phase and specific inhibition of 28S and 5.8S rRNA synthesis, as well as a deficit in the
	cytosolic 60S ribosomal subunit. This suggests that BOP1 is involved in the formation of
	mature rRNAs and in the biogenesis of the 60S ribosomal subunit. BOP1 physically interacts
	with pescadillo (a protein involved in cell proliferation) and enables efficient incorporation of
	pescadillo into the nucleolar preribosomal complexes, thereby affecting rRNA maturation and
	the cell cycle. The BOP1-pescadillo complex is also necessary for biogenesis of 60S ribosomal
	subunits. Deregulation of BOP1 may lead to colorectal tumorigenesis.
Gene ID:	23246
Application Details	
Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	
Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative: ProClin	

## Handling

Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months