

[Go to Product page](#)

## Datasheet for ABIN1714140 **anti-Cyclin Y antibody (AA 21-120)**

### Overview

Quantity:	100 µL
Target:	Cyclin Y (CCNY)
Binding Specificity:	AA 21-120
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This Cyclin Y antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin- embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

### Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Cyclin Y/Cyclin X
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Dog,Cow,Sheep,Pig,Chicken,Rabbit
Purification:	Purified by Protein A.

### Target Details

Target:	Cyclin Y (CCNY)
Alternative Name:	Cyclin Y/Cyclin X ( <a href="#">CCNY Products</a> )

## Target Details

Background:	<p>Synonyms: C10orf9, CBCP1, CCNX, CCNY, CFP1, Chromosome 10 open reading frame 9, Cyclin box carrying protein 1, Cyclin box protein 1, Cyclin fold protein 1, Cyclin X, Cyclin Y, CCNY_HUMAN.</p> <p>Background: Cyclin Y is a 341 amino acid protein belonging to the cyclin family. Cyclin Y exists as three alternatively spliced isoforms and contains a cyclin N-terminal domain. Cyclin Y may control cell division cycles and regulate cyclin-dependent kinases. Cell proliferation is controlled at specific stages of the cell cycle by distinct protein kinase complexes. These complexes consist of a catalytic subunit associating with a specific regulatory subunit to form the active kinase. The cyclins, which include cyclin A, B, C, D, E, F, G, H, I, K, L, T, Y and their related proteins, including Dbf4, comprise the regulatory subunits of these kinase complexes. The controlled activation of the kinase complexes at various intervals of the cell cycle is regulated by the availability of the cyclins to the catalytic subunit. Unlike the catalytic subunit, which is expressed continually, the expression and stability of the regulatory subunit fluctuates depending on the stage of the cell cycle, thereby regulating kinase activity.</p>
-------------	--

Gene ID:	219771
----------	--------

## Application Details

Application Notes:	WB 1:300-5000 ELISA 1:500-1000 IHC-P 1:200-400 IHC-F 1:100-500 IF(IHC-P) 1:50-200 IF(IHC-F) 1:50-200 IF(ICC) 1:50-200 ICC 1:100-500
--------------------	--

Restrictions:	For Research Use only
---------------	-----------------------

## Handling

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS( pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

---

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months