antibodies

Datasheet for ABIN1714192 anti-GC-Rich Promoter Binding Protein 1 (GPBP1) (AA 11-110) antibody



Overview

Quantity:	100 µL
Target:	GC-Rich Promoter Binding Protein 1 (GPBP1)
Binding Specificity:	AA 11-110
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	Un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin- embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human GPBP1/Vasculin
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Dog,Cow,Sheep,Pig,Horse,Chicken
Purification:	Purified by Protein A.

Target Details

Target:

GC-Rich Promoter Binding Protein 1 (GPBP1)

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 1/3 | Product datasheet for ABIN1714192 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

Target Details	
Alternative Name:	GPBP1/Vasculin (GPBP1 Products)
Background:	Synonyms: DKFZp761C169, GC rich promoter binding protein 1, GC-rich promoter-binding
	protein 1, GPBP, Gpbp1, GPBP1_HUMAN, MGC126339, SSH6, Vascular wall linked, Vascular
	wall-linked protein, Vasculin.
	Background: Vascular wall-linked protein, or Vasculin, is primarily expressed in the arterial wall
	and in plasma. It is also differentially expressed in human atherogenesis. Alternative splicing of
	exon 3 of the Vasculin gene produces three variants. Vasculin binds to and activates the
	minimal self-sufficient promoter element (MSPE) of the mouse Ada gene promoter and binds
	to and partially suppresses the GC-rich promoter of the nonhomologous human TOP2A gene
	promoter. It acts as a nuclear factor that can form complexes with TATA-binding proteins,
	transcription factors TFIIB and TFIIF, RNA polymerase II and p300. The regulated expression of
	Vasculin in plaques suggests that it may be involved in atherogenesis, and its presence in
	plasma may implicate Vasculin as a marker for atherosclerosis.
Gene ID:	65056

Application Details

Application Notes:	WB 1:300-5000
	ELISA 1:500-1000
	IHC-P 1:200-400
	IHC-F 1:100-500
	IF(IHC-P) 1:50-200
	IF(IHC-F) 1:50-200
	IF(ICC) 1:50-200
	ICC 1:100-500
Restrictions:	For Research Use only
Handling	

Format:	Liquid
Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Order at www.antibodies-online.com | www.antikoerper-online.de | www.anticorps-enligne.fr | www.antibodies-online.cn International: +49 (0)241 95 163 153 | USA & Canada: +1 877 302 8632 | support@antibodies-online.com Page 2/3 | Product datasheet for ABIN1714192 | 03/08/2024 | Copyright antibodies-online. All rights reserved.

Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months