

Datasheet for ABIN1714302
anti-ARID3A antibody (AA 201-300)



[Go to Product page](#)

Overview

Quantity:	100 µL
Target:	ARID3A
Binding Specificity:	AA 201-300
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This ARID3A antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human DRIL1/ARID3A
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Cow, Sheep, Pig, Horse, Chicken
Purification:	Purified by Protein A.

Target Details

Target:	ARID3A
Alternative Name:	DRIL1/ARID3A (ARID3A Products)

Target Details

Background: Synonyms: AR13A_HUMAN, ARID domain-containing protein 3A, ARID3A, AT rich interactive domain-containing protein 3A, AT-rich interactive domain-containing protein 3A, B cell regulator of IgH transcription, B-cell regulator of IgH transcription, Bright, dead ringer like 1, Dead ringer-like protein 1, DRIL1, DRIL3, E2F binding protein 1, E2F-binding protein 1, E2FBP1.

Background: ARID3A, also known as DRIL1 in humans and Bright (for B cell regulator of IgH transcription) in mice, are the mammalian homologs of the Drosophila Dri (dead ringer) protein. ARID3A is developmentally regulated and is expressed in a restricted set of cells, including differentiating cells of the gut and salivary glands. ARID3A represents a member of a unique family of transcriptional activators that shares sequence similarity to proteins of SWI/SNF complexes, it contains an A/T-rich DNA-binding (ARID) domain and a distinct domain involved in tetramerization. The gene encoding ARID3A is linked to a marker of Peutz-Jeghers syndrome, which is an autosomal-dominant disorder characterized by melanocytic macules of the lips, multiple gastrointestinal hamartomatous polyps and an increased risk for various neoplasms, including gastrointestinal cancer. E2FBP1 (E2F-1 binding protein 1) is identical to ARID3A in the carboxy terminal region. E2FBP1 appears to lack DNA binding and transactivation domains, and it functions to regulate the transcription of proteins involved in cell proliferation by binding to the transcription factor E2F-1.

Gene ID: 1820

Pathways: [Chromatin Binding](#)

Application Details

Application Notes: WB 1:300-5000
ELISA 1:500-1000
IHC-P 1:200-400
IHC-F 1:100-500
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
ICC 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Handling

Concentration:	1 µg/µL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months