

[Go to Product page](#)

Datasheet for ABIN1714306 **anti-TXNDC12 antibody (AA 21-120)**

Overview

Quantity:	100 µL
Target:	TXNDC12
Binding Specificity:	AA 21-120
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This TXNDC12 antibody is un-conjugated
Application:	ELISA, Western Blotting (WB), Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin- embedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human ERp19
Isotype:	IgG
Predicted Reactivity:	Human, Mouse, Rat, Cow, Sheep, Pig, Horse, Chicken
Purification:	Purified by Protein A.

Target Details

Target:	TXNDC12
Alternative Name:	ERp19 (TXNDC12 Products)

Target Details

Background: Synonyms: AG1, AGR1, anterior gradient homolog 1, endoplasmic reticulum protein ERp19, endoplasmic reticulum resident protein 18, endoplasmic reticulum resident protein 19, endoplasmic reticulum thioredoxin superfamily member, 18 kDa, ER protein 18, ER protein 19, ERP 18, ERP16, ERP19, hAG 1, hAG1, hTLP19, PDIA16, protein disulfide isomerase family A, member 16, thioredoxin domain containing 12 endoplasmic reticulum, Thioredoxin domain-containing protein 12, thioredoxin like protein p19, TLP19, TXNDC12, TXD12_HUMAN.

Background: Endoplasmic reticulum proteins (ERps) are widely expressed proteins and localize to the ER. ERp19, ERp29, ERp46, ERp57 and ERp72 may act as proteases, protein disulfide isomerases, thiol-disulfide oxidases, phospholipases or a combination of these. ERp19, also designated thioredoxin domain-containing protein 12 (TXNDC12), and ERp46, also designated thioredoxin domain containing 5 (TXNDC5), belong to the thioredoxin superfamily and contain a thioredoxin fold with a consensus active-site sequence (CxxC). Both ERp19 and ERp46 are widely expressed ER luminal proteins that are most abundant in the liver and are enriched in purified liver ER vesicles. ERp19 shows significant protein thiol-disulfide oxidase activity in vitro, which is dependent on the presence of both active-site cysteines.

Gene ID: 51060

Pathways: [Cell RedoxHomeostasis](#)

Application Details

Application Notes: WB 1:300-5000
ELISA 1:500-1000
IHC-P 1:200-400
IHC-F 1:100-500
IF(IHC-P) 1:50-200
IF(IHC-F) 1:50-200
IF(ICC) 1:50-200
ICC 1:100-500

Restrictions: For Research Use only

Handling

Format: Liquid

Concentration: 1 µg/µL

Buffer: 0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.

Handling

Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months