antibodies -online.com







Go to Product page

Overview

Quantity:	100 μL
Target:	FLRT1
Binding Specificity:	AA 301-400
Reactivity:	Human
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This FLRT1 antibody is un-conjugated
Application:	Western Blotting (WB), ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffin-embedded Sections) (IF (p)), Immunohistochemistry (Paraffin-embedded Sections) (IHC (p)), Immunocytochemistry (ICC), Immunohistochemistry (Frozen Sections) (IHC (fro))

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human FLRT1
Isotype:	IgG
Predicted Reactivity:	Human,Mouse,Rat,Cow,Sheep,Pig,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	FLRT1
Alternative Name:	FLRT1 (FLRT1 Products)

Target Details

Bacl	//	ro	ıın	٨.
Daci	ĸα	ΙU	uu	u.

Synonyms: Fibronectin-like domain containing leucine rich transmembrane protein 1;
Fibronectin-like domain-containing leucine-rich transmembrane protein 1; FLRT1;

FLRT1_HUMAN; Leucine rich repeat transmembrane protein FLRT1; Leucine-rich repeat

transmembrane protein FLRT1.

Background: The leucine-rich (LRR) repeat is a 20-30 amino acid motif that forms a hydrophobic å/ horseshoe fold, allowing it to accommodate several leucine residues within a tightly packed core. All LRR repeats contain a variable segment and a highly conserved segment, the latter of which accounts for 11 or 12 residues of the entire LRR motif. FLRT1 (fibronectin leucine rich transmembrane protein 1) is a 646 amino acid single-pass type I membrane protein that contains one fibronectin type-III domain and ten LRR repeats. Expressed in kidney and brain, FLRT1 is thought to play a role in cell adhesion and receptor signaling. FLRT1 shares similarity with FLRT2 and FLRT3 and is subject to post-translational N-glycosylation. The gene encoding FLRT1 maps to human chromosome 11, which houses over 1,400 genes and comprises nearly 4% of the human genome.

Gene ID:

23769

Application Details

Application Notes:

WB 1:300-5000

ELISA 1:500-1000

IHC-P 1:200-400

IHC-F 1:100-500

IF(IHC-P) 1:50-200

IF(IHC-F) 1:50-200

IF(ICC) 1:50-200

ICC 1:100-500

Restrictions:

For Research Use only

Handling

Format:	Liquid
Concentration:	1 μg/μL
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.
Preservative:	ProClin
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be

Handling

	handled by trained staff only.
Storage:	4 °C,-20 °C
Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.
Expiry Date:	12 months