antibodies -online.com







anti-LIMD1 antibody (AA 21-120)



Overview

Quantity:	100 μL
Target:	LIMD1
Binding Specificity:	AA 21-120
Reactivity:	Mouse
Host:	Rabbit
Clonality:	Polyclonal
Conjugate:	This LIMD1 antibody is un-conjugated
Application:	ELISA, Immunofluorescence (Cultured Cells) (IF (cc)), Immunofluorescence (Paraffinembedded Sections) (IF (p)), Immunohistochemistry (Paraffinembedded Sections) (IHC (p)), Immunohistochemistry (Frozen Sections) (IHC (fro)), Immunocytochemistry (ICC)

Product Details

Immunogen:	KLH conjugated synthetic peptide derived from human Limd1
Isotype:	IgG
Cross-Reactivity:	Mouse
Predicted Reactivity:	Human,Rat,Cow,Sheep,Pig,Chicken,Rabbit
Purification:	Purified by Protein A.

Target Details

Target:	LIMD1

Target Details

Target Details		
Alternative Name:	Limd1 (LIMD1 Products)	
Background:	Synonyms: AW822033, D9Ertd192e, LIM domain-containing protein 1, LIM domains containing	
	1, LIMD1, LIMD1_HUMAN, OTTHUMP00000164657, OTTHUMP00000210317, rCG_25194.	
	Background: The Zyxin family of proteins contains five members: Ajuba, Limd1, LPP, TRIP6 and	
	Zyxin. Limd1 (LIM domain-containing protein 1) is a ubiquitously expressed tumor suppressor	
	containing 3 LIM zinc-binding domains. LIM domains consist of a cysteine-rich consensus	
	sequence containing two distinct zinc-binding subdomains, which mediate protein-protein	
	interactions. Limd1 interacts with the proteins SQSTM1, Rb, p62 and TRAF6. Limd1 was first	
	identified when the deletion of its gene was noted in some cervical cancers. Limd1 blocks in	
	vitro and in vivo tumor growth and is down-regulated in lung cancer. Limd1 may regulate	
	osteoclast development under stressful conditions via its interactions with TRAF6 and p62.	
Gene ID:	8994	
Pathways:	Ribonucleoprotein Complex Subunit Organization	
Application Details		
Application Notes:	ELISA 1:500-1000	
	IHC-P 1:200-400	
	IHC-F 1:100-500	
	IF(IHC-P) 1:50-200	
	IF(IHC-F) 1:50-200	
	IF(ICC) 1:50-200	
	ICC 1:100-500	
Restrictions:	For Research Use only	
Handling		
Format:	Liquid	
Concentration:	1 μg/μL	
Buffer:	0.01M TBS(pH 7.4) with 1 % BSA, 0.02 % Proclin300 and 50 % Glycerol.	
Preservative:	ProClin	
Precaution of Use:	This product contains ProClin: a POISONOUS AND HAZARDOUS SUBSTANCE, which should be handled by trained staff only.	
Storage:	4 °C,-20 °C	

Handling

Storage Comment:	Shipped at 4°C. Store at -20°C for one year. Avoid repeated freeze/thaw cycles.	
Expiry Date:	12 months	